



## LJMU Research Online

**Premathilake, R and Hunt, CO**

**Late Pleistocene humans in Sri Lanka used plant resources: A phytolith record from Fahien rock shelter**

<http://researchonline.ljmu.ac.uk/id/eprint/8819/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Premathilake, R and Hunt, CO (2018) Late Pleistocene humans in Sri Lanka used plant resources: A phytolith record from Fahien rock shelter. *Palaeogeography, Palaeoclimatology, Palaeoecology*. ISSN 0031-0182**

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

**Late Pleistocene humans in Sri Lanka used plant resources: a phytolith record from Fahien Rock Shelter**

Rathnasiri Premathilake<sup>1</sup>, Chris O. Hunt<sup>2</sup>

<sup>1</sup>Postgraduate Institute of Archaeology, University of Kelaniya, Sri Lanka (Email: [premathilake@hotmail.com](mailto:premathilake@hotmail.com))

<sup>2</sup>School of Natural Sciences and Psychology, Liverpool John Moores University UK (Email: [C.O.Hunt@ljmu.ac.uk](mailto:C.O.Hunt@ljmu.ac.uk))

## **Abstract**

Little is known of the human use of rainforest plant resources of prehistoric Sri Lanka due to the lack of preservation of organic material and the effects of various destructive taphonomic processes. Phytoliths recovered from a AMS radiocarbon and OSL dated sequence at Fahien Rock Shelter indicate interactions of anatomically modern humans with the lowland rainforests of south-western Sri Lanka from 44,952-47,854 cal. BP to 11,991-12,402 cal. BP. During this period, the Rock Shelter occupants extracted their livelihood from a number of wild plants including bananas, rice, breadfruits, durians, canarium and species of palm and bamboo. These taxa are associated with present-day disturbed lowland rainforests. Gathering and processing of plant resources by existing modern rainforest foragers cannot directly be compared with the subsistence activities of the Late Pleistocene Rock Shelter occupants.

Key words: Excavation; Stratigraphy; Taphonomy; Rainforests; Rice; Wild Banana

## 1. Introduction

Tropical environments are reputed to be amongst the most challenging for human colonisation (Friesem et al., 2016). Until recently, many anthropologists held that it was impossible for hunter-gatherers to live in rainforest environments if they could not exchange forest resources with adjacent agricultural populations (Headland and Bailey, 1991). Ethnohistorical, geomorphological, palaeoecological, archaeological and archaeobotanical evidence is important to examine questions concerning the antiquity of rainforest colonization by humans on a global scale (Cosgrove et al., 2007; Barker et al., 2007, 2017). However, there is emerging archaeobotanical evidence (e.g. phytoliths, pollen and macro-plant remains) to indicate that humans were able to colonise and exploit rainforest environments in Island South East Asia, Melanesia, Australasia, Sri Lanka and Africa from the late Pleistocene and in doing so, they utilised fire to manage rainforest vegetation (e.g. Premathilake, 2003, 2006; Barker et al., 2007, 2017; Petraglia et al., 2009; Clarkson et al., 2009; Lentfer et al., 2010; Summerhayes et al., 2010; Perera, 2010; Perera et al., 2011; Hunt et al., 2012; Castillo and Fuller, 2015; Roberts and Petraglia, 2015; Roberts et al., 2015, 2017).

Phytoliths - minute silica bodies produced by plants (Piperno, 1988) - are one of the most promising lines of evidence in tropical environments. These environments are often highly oxidizing, with frequent free iron, aluminium oxides and other clay minerals and thus with poor preservation of organic materials such as pollen, plant macrofossils and wood. This has major implications in reporting the history of use of rainforest plant resources (Deraniyagala, 1992; Goldberg and Bar-Yosef, 1998; Bowdery, 1999; Albert et al., 1999; Denham et al., 2003; Premathilake, 2003, 2006; Fuller and Harvey 2006; Vrydaghs et al., 2009; Castillo and Fuller, 2010; Perera 2010; Perera et al., 2011; Madella and Lancelotti, 2012; Castillo and Fuller, 2015; Friesem et al., 2016; Bates et al., 2017; Premathilake et al., 2017a,b). We report the results of phytolith analysis from the deposits at Fahien Rock Shelter, one of the oldest prehistoric sites in South Asia, which contains a sequence containing evidence for habitation by rainforest hunter-gatherers through the second half of the late Pleistocene and into the Holocene.

## 2. Fahien Rock Shelter environment

Fahien Rock Shelter, one of the largest caves in Sri Lanka, is situated at 6° 38' 55" N 80° 12' 55" E and 130 m above sea-level close to Yatagampitiya village, near Bulathsinhala in the Kalutara District, southwest Sri Lanka (Fig. 1). It is a complex of interconnected Rock Shelters developed in coarse crystalline gneiss rock faces (Cooray, 1984). The mouth has a width of 30 m and average height above the floor of 20 m. The interior is about 10 m deep and slopes down west to east. Data from the meteorological station in Sinharaja, in lowland rainforest some 10 km away (06°23'22"N 80°30'05"E) indicate that the regional climate is humid-tropical with a mean annual temperature of 26°C and annual average rainfall of 4300 mm (Zoysa and Raheem, 1987; Gunatillake et al., 2004). Today, the landscape around the site is characterized by disturbed lowland rainforest, with paddy fields present in the slightly incised valley below the Rock Shelter.

## 3. Material and methods

The Rock Shelter was excavated over several seasons between 1986 and 2012 by scholars from the Department of Archaeological Survey, Sri Lanka (Deraniyagala, 1992; Wijeyapala, 1997; Perera, 2010, 2015; Oshan, 2011). The excavation area (4 x 5 m) is located in the east of the main chamber of the Rock Shelter, and a detailed stratigraphical study was undertaken. The depiction of the stratigraphic succession with its contained archaeological contexts at the Rock Shelter was made according to the standard Harris Matrix tool (cf. Premathilake and Hunt, 2018).

### 3.1. Lithostratigraphy

The excavation penetrated ca. 2.40 m of heterogeneous clast-rich loamy sediments which can be subdivided into five major stratigraphic units, 10 archaeological phases (I-X) and approximately 250 archaeological contexts (C). This paper considers the lithological records from units L5 and L4/L3, including phases I-VI of the archaeological sequence, where the most significant late Pleistocene archaeological evidence was found. This includes the oldest microlith toolkits known to South Asia, associated with the contexts from C-92 to C-87 (Fig. 2; Table 1).

### 3.2. Chronostratigraphy

Nine AMS dates were obtained from charred materials, charred wood, charcoal, *Canarium cf. zylanicum* nut and freshwater shells deposited throughout the investigated southern section. Two sediment samples (SUTL2327, SUTL2326) from context C-92 were used for OSL dating. Radiocarbon dates were produced using Accelerator Mass Spectrometry at the CHRONO Centre at Queen's University, Belfast and the Beta Analytic Laboratory in the USA. The dates were calibrated using Calib 6.11 (Reimer et al., 2009). The radiocarbon dates indicate that the studied sediments were laid down between 44,950-47,854 cal. BP and 11,991-12,402 cal. BP (Table 2; Fig. 2). The oldest microlith toolkits known to South Asia are associated with the contexts from C-92 to C-87, which are dated to between 44,950-47,854 cal. BP and 27,860-29,089 cal. BP. OSL dating (Kinnaird & Sanderson, 2010) appears to be broadly consistent with the radiocarbon chronology (Premathilake and Hunt, 2018).

### 3.3. Biostratigraphy

Table 1 includes macro-botanical and zoological remains reported through the stratigraphy. This comprises shell beads, animal teeth (e.g. *Semnopithecus priam*, *Ratufa macroura*, *Parradoxurus* sp., *Trachypithecus vetulus* and *Macaca sinica*), human bones, burnt and unburnt shells (e.g. *Acavus* sp. and *Pila* sp.), charcoal, wild breadfruit (*Artocarpus nobilis*), epicarps, *Canarium* nuts and coprolites. Human remains include several internments; some coated with red ochre and are associated with the first microlith and osseous technologies anywhere in South Asia. The layer is dated on charcoal and charred materials to 39,852-35,768 cal. BP (Table 2). Phase VI in the sequence has been characterised as a period of intensive human habitation at the site, as indicated by the presence of a highly fragile and fragmentary human skeleton (Perera, 2010, 2015; Perera et al., 2011). This has recently been dated to around 12,000 cal. BP. (cf. Perera, 2015). The stratigraphy also contains well-preserved ashy hearths.

### 3.4. Sample selection and sediment processing

Twelve 30x10x8 cm monoliths were taken from the southern profile of the Rock Shelter. These covered five major layers (L3-L5) including archaeological phases I-VI; (Table 1) (Wijeyapala, 1997; Perera, 2010, 2015; Oshan, 2011). Six subsamples (C-92 - C-81) from

the lowermost sediments of late Pleistocene age including L5 and L4/L3 were selected for detailed phytolith analysis on the basis of their archaeological and chronological significance (cf. Premathilake and Hunt, 2018). The detailed techniques used for phytolith extraction (Lentfer and Boyd, 1998), microscopy and phytolith percentage diagram construction are presented in SOM 1.

### 3.5. Phytolith taxonomy

Classification and taxonomic identifications of phytoliths from archaeological samples were made using modern and archaeological phytolith collections housed at the Laboratory for Palaeoecology, Postgraduate Institute of Archaeology, University of Kelaniya, Sri Lanka and the French Institute of Pondicherry, India (IFP). In our procedures, the most common morphological characteristics, general occurrence and appearance of key archaeological phytoliths were comparatively studied (Figs. 3, 4, 5, 6). Additional support received from several published works and details of taxonomic separation are presented in SOM 1 and 2.

## 4. Results

### 4.1. Phytoliths stratigraphy

Phytolith assemblages are shown in Fig. 7 and Table S3. All samples contained high counts of well-preserved phytoliths. In a very few samples, some phytoliths are pitted, displayed a few relatively large micro-channels, mineralized micro-structures or were broken. More than 60 phytolith morphotypes were identified. Monocotyledonous phytoliths were dominant in comparison to the dicotyledonous types. The detailed results are given in Table S3. Phytolith counts decrease from the Late to Terminal Pleistocene samples.

Phytoliths from seeds and leaves of wild *Musa* banana commonly occur (SOM 2; Table S3). Phytoliths from several other economically-important taxa e.g. species of *Palmae*, *Artocarpus* cf. *nobilis* (wild breadfruit) and *Durio* occur in relatively low percentages. One terminal Pleistocene sample contained a few phytoliths of *Burseraceae*. The presence of phytoliths from *Oryza* sp. (wild rice) seeds and leaves is reported in varying percentages (SOM 2; Table S3). Phytoliths from *Poaceae* (grass) occur in all samples, with Pooid regular short cells relatively common compared with Chloridoid saddles and Panicoid bilobates. All samples contain bulliforms derived from *Poaceae*. Rondel phytoliths are common in all the samples.

Many of them are distinctly taller than width (corn-like rondel), most possibly derived from Bambusoid species. A few samples contain Cyperaceae phytoliths.

Phytoliths blackened by fire (reported as burnt phytoliths) are common. Many of the burnt phytoliths are from Poaceae, Palmae (identified as Palmae 1 and 2 based on size variation), *Oryza* sp., *Durio* sp. and *Artocarpus* cf. *nobilis*.

Freshwater diatom (*Navicula* cf. *leistikowii* and *Aulacoseria* cf. *granulata*) counts increase from the Late to the terminal Pleistocene samples. A few brackish/marine diatoms (*Navicula* cf. *gregaria*, *Amphora* cf. *coffeaeformis* and *Achnanthes* cf. *hauckiana*) occur in several samples (Table S3).

## 5. Discussion

### 5.1. Reliability of Rock Shelter stratigraphy

Understanding of the litho-stratigraphy is imperative for interpreting the Rock Shelter phytoliths, although many challenges still lies ahead of investigations of how sediment stratigraphies can develop (Gilbertson et al. 2005; Butzer, 2008; Hunt et al., 2010; Friesem, 2016; Zurro et al., 2016; Barker et al, 2007, 2017). The sediments of Fahien Rock Shelter are heterogeneous and analysis of the ca. 250 contexts suggests the occurrence of very complex sedimentary processes (Perera, 2010, 2015; Kourampas et al., 2009; Oshan, 2011). Nevertheless, all of the radiocarbon dates are in good stratigraphic order, suggesting sequential deposition and little post-depositional disturbance.

The chronology suggests that significant depositional hiatuses occur within the excavated sequence between the late Pleistocene and early Holocene (Fig. 2; Table 2). The oldest sample, from context 92; just above the date of 44,952-47,854 cal BP contains a very high number of phytoliths suggesting significant human activity (Premathilake and Hunt, 2017). A major hiatus extended from C-87 to C-81 (27,860-29,089 to 11,991-12,402cal. BP) and is marked by the reduction of phytolith sums (Fig. 7).

These hiatuses can be explained by climatic phases causing erosion of the Rock Shelter sediments. Multi-proxy investigations (Premathilake and Risberg, 2003; Petraglia et al., 2009; Premathilake, 2006, 2012ab, 2015a) indicate Inter Tropical Convergence Zone (ITCZ) -



induced monsoon intensification events with high rainfall at 17,650-16,200, 13,700-12,800, 10,400-9,900 and 9,200-8,800 cal. BP, contemporary with periods of increased rainfall-driven landslide frequency in the landscape around the Rock Shelter (Gunathilaka, 2007; Kourampas et al., 2009; Premathilake and Hunt, 2018). These alternate with desiccation from millennial to multi-century scale arid/semi-arid phases of monsoon failure, identified from peat and sedimentary archives in southern Asia between 24,000 cal. BP and 8,100 cal. BP (Premathilake and Risberg, 2003; Premathilake and Gunathilaka, 2013). Phases of erosion of the Rock Shelter sediments (Fig. 2) seem to be contemporary with the high-rainfall episodes, suggesting that the impact of climate was the dominant factor forming the litho-stratigraphy throughout the late Pleistocene. We argue that accumulation of clast-rich loam sediment layers (Table 1) continued because of the combined effect of the prominent mode of semi-arid climate and human activities including biomass burning and foraging associated with lowland and upper montane rain forests (Perera et al., 2011, Kourampas et al., 2009, Premathilake, 2003, 2006). It seems that anthropogenic sedimentation, particularly accumulation of human habitation debris and sediment entrained by human action was prominent during the late Pleistocene, but the accumulation rate varied very markedly between and within major layers through this time (Fig. 2). Detailed geo-archaeological research (soil micromorphology) with additional radiocarbon dates will provide more information that will further enhance and clarify the current understanding of the depositional history of the sequence (Friesem et al., 2016).

## *5.2. Reliability of phytolith assemblages*

Seeds, epicarps, charcoal and wood recovered from several rock shelter deposits in south western Sri Lanka have been used to address archaeological issues including subsistence activities of the rock shelter occupants and understanding site environments but, phytoliths have not been discussed in detail. Understanding the late Pleistocene rock shelter sites is often difficult because of poor preservation of organic materials and the effect of various taphonomic processes (Deraniyagala, 1992; Perera, 2010; Perera et al., 2011; Kourampas et al., 2009; Roberts et al., 2015, 2017). In the case of Fahien Rock Shelter, taphonomic investigation was essential to understand how phytoliths reached the site and how they were deposited and preserved. The presence of phytoliths in the Rock Shelter sediments provides information about the depositional processes in several ways (1) *in situ* plant decay leading to phytolith

deposition on surfaces (2) alluvial or colluvial re-deposition of phytoliths along with their associated sediments (3) wind deposition (4) cultural deposition of phytoliths through plant materials used by Rock Shelter occupants for food and other cultural purposes (Behrensmeyer et al., 2000; Hunt et al., 2016).

The lack of macro botanical evidence – fruits, leaves, stems etc. - of living plants of taxa such as Poaceae, Palmae and Musaceae and the lack of organic litter in the Rock Shelter suggest a minimal deposition of *in situ* phytoliths. Abundant phytoliths from these monocot taxa in the samples suggest that the effect of horizontal and vertical movements through alluvial and/or colluvial processes may have played a limited role in phytolith re-deposition. Due to the presence of a drip-line, rain water penetration into the Rock Shelter is minimal. Wind deposition is rare due to the particular geomorphology of Rock shelter in the humid tropical environment.

The main process, therefore, of phytolith deposition in the Rock shelter is most likely to have been through human or animal vectors. Common animals in the vicinity of the Rock Shelter, such as bats, birds, and insects, are rather unlikely to play a role in phytolith deposition, however. The variable phytolith counts through the sequence (Fig. 7) and selective distribution of phytoliths from wild bananas suggest that most phytoliths were brought into the cave by people and remain *in-situ* (cf. Premathilake and Hunt, 2018). Hunt et al. (2016) suggested that similar strong patterning at Niah Cave was evidence for anthropogenic deposition. Thus, it is inferred that humans are the most likely agents for phytolith deposition – with materials from economically significant plants such as wild banana, breadfruit and palms from plants characteristically growing in disturbed lowland rainforest near to the Rock Shelter (most likely within a few kilometers at most). Abundant phytoliths from grasses and sedges - are identified as from economically-useful taxa, but may also be evidence for forest disturbance. The occurrence of freshwater and brackish-marine diatom species throughout is not surprising in habitation deposits (Hunt et al. 2016) and is consistent with a number of human activities involving activity in wetlands and the carrying of material encrusted with diatoms into the Rock Shelter (Kanthilatha et al., 2017).

In the majority of samples, abundant banana seed and leaf phytoliths together with the lack of taphonomic markers (such as breakage, corrosion, microchannels, regulation, dissolution pits, mineralized microstructures, cut marks and pitted patterns) indicate excellent preservation conditions and selective distribution of phytoliths from wild bananas used by the Rock Shelter occupants. This suggests high phytolith compositional fidelity in the samples. Excellent preservation conditions are further attested by the high number of non-durable phytoliths (e.g. globular echinate, volcaniforms, trichome, elongate sinuous and bilobate) and very few phytoliths with isolated cavities, with occluded carbon. A few phytoliths contain cavities that may be due to the natural silicification process, as evidenced by the modern phytolith collections and are not related to taphonomic processes such as diagenetic dissolution. The presence of well-preserved phytoliths throughout the sequence suggests these impacts were minimal (Alexandre et al., 1997; Behrensmeyer et al., 2000; Fredlund and Tieszen, 1997; Albert et al., 1999, 2006; Farmer et al., 2005; Piperno, 2006; Gérard, et al., 2008; Loucaides et al., 2008, 2010; Osterrieth, et al., 2009; Karkanas, 2010; Borrelli et al., 2010; Cabanes et al., 2011, 2012; Alexandre et al., 2015). Alkaline conditions are thought to contribute to phytolith dissolution (Loucaides et al., 2010; Cabanes et al., 2011; Fraysse et al., 2006, 2009) with an increase in solubility of silica at  $\text{pH} > 7.8$ . In the iron (Fe) rich fine-grained Fahien Rock Shelter sediments solubility is limited as indicated by pH measurements of 6.5-7.3 in all the sediment samples studied (cf. Piperno, 1988). Facetate and sclereid phytoliths from woody dicotyledonous (e.g. forest taxa) are rare in the Late Pleistocene and Holocene samples, which might possibly due to dissolution (Piperno, 2006), but given the circumneutral sediments it is more likely that few phytoliths from woody materials were incorporated. We acknowledge that the production of facetates and sclereids from modern forest plant species is relatively high, but they are very unlikely to be preserved in much older samples (Premathilake et al., 2017b).

Except in the very uppermost parts of the sequence, the absence of post-depositional disruption through root penetration, roof fall, vertical sediment cracks, human digging and animal burrowing and the lack of evidence for disruption within the clay- and silt-rich, highly-compacted and multi-layered sediments indicates likely very limited vertical movement of phytoliths (Perera, 2010; 2015). The distribution of the smallest phytoliths (e.g. small rectangles

from wild banana seeds and globular nodules from Bombacaceae) and fine-grained sediments suggests the minimal impact of illuviation of clay minerals, reported as a common process in some rock shelter stratigraphies (Butzer 2008; Kourampas, 2009; Alexandre et al., 1997). All these minimized sources of bias indicate that spatial and temporal fidelity is relatively high in this Rock Shelter phytolith stratigraphy (Behrensmeyer et al., 2000). Therefore phytoliths buried in deep stratigraphic levels are likely to be securely *in-situ* and unaffected by bioturbation, suffusion, infiltration or corrosion (Premathilake and Hunt, 2018).

### 5.3. Late Pleistocene plant exploitation

The Rock Shelter phytolith assemblages provide a potentially more complete botanical record than that provided by macrofossils (Table 1). The significant phytolith counts of economically important plants - wild bananas (*Musa accuminata* and *Musa balbisiana*), wild rice (*Oryza* spp), breadfruit (*Artocarpus* cf. *nobilis*), *Durio*, sp. Palmae and grasses - suggests that the Rock Shelter occupants used those plants. Significant late Pleistocene archaeological evidence is associated with contexts C-87 to C-92; which are dated to between 44,950-47,854 and 27,860-29,089 cal. BP. The deposits in those contexts yielded high counts of phytoliths from economic plants. Key taxa are discussed below.

#### 5.3.1. Wild bananas

Today, more than 50 *Musa* species are recorded in tropical and subtropical regions (Cheesman, 1947; Simmonds, 1956, 1962; Constantine, 1999-2008; Fuller and Madella, 2009; Häkkinen and Väre, 2008). In Sri Lanka, the perennial species *M. acuminata* and *M. balbisiana* commonly grow in disturbed lowland rainforest from sea-level up to 900 m, and they are used for food and various cultural purposes by local people (Chandraratne and Nanayakkara, 1951; Perera, 2017). The Pali chronicle '*Mahavamsa*', written in the 5<sup>th</sup> century A.D., reported wild banana (*attikehel*) with its various uses. Ethnobotanical investigations indicate that immature and mature fruits with or without seeds, leaves, young leaf sheaths, the male inflorescence bud, male flowers, fruit peels, pseudostem, sucker/shoots, rhizome and ash of wild banana were widely used for plates, wrapping foods, medicine, stimulants, textiles, clothing, packaging, paper-making, craft industry, ornaments and also ceremonial, magic and ritual activities in the

countries of south and southeast Asia (Christiansen, 2002; Fuller and Madella, 2009; Simmonds, 1956, 1962; Kennedy, 2009).

High counts of wild *Musa* banana seed and leaf phytoliths (Table S1, S2, S3) indicate that the Rock Shelter occupants exploited wild bananas throughout the late Pleistocene. The banana phytoliths in the late Pleistocene samples are identical to those found in modern *M. accuminata* and *M. balbisiana* populations in Sri Lanka (Premathilake, 2016; Perera, 2017; Premathilake and Hunt, 2018; SOM 2; Fig. 8). This suggests that these two species existed as natural populations in association with open semi-natural habitats in heavily human-impacted lowland rainforest of south western Sri Lanka (Fuller and Madella, 2009; De Langhe, 2009; Roberts et al., 2015, 2017). The suggestion of open habitats is consistent with the abundant wild banana seeds and scarcity of woody dicotyledonous phytoliths at Fahien (Fig. 7). The antiquity of wild banana use, previously known from charred seeds from the terminal Pleistocene at Beli-lena rock shelter (Kajale, 1989), has been pushed back into the late Pleistocene and the date of 44,950-47,854 cal. BP by this study. The variety of wild banana phytoliths at Fahien is consistent with multiple uses including the fruits for starchy food, leaves for ethnographically-attested uses such as plates and clothing, stems for fuel. The common occurrence of burnt (blackened) banana seed phytoliths associated with ashy deposits, charcoal and bone fragments from hearths (Perera, 2010; Fig. 7; Table 1; Table S3); suggests perhaps that the roasting and inadvertent burning of banana fruits may have released blackened phytoliths, suggesting possible employment of food processing techniques (Heyne 1950; Perera, 2010; Kanthilatha et al., 2017). However, the available phytolith evidence is not enough to understand the relationship between the phytolith assemblages yielded by processing techniques and the phytoliths found in Rock Shelter floors (Kanthilatha et al., 2017). It is possible given the evidence for behavioural richness of the occupants, including the use of shark vertebra, marine shells as beads and a shell pendant at 38,000 cal. BP (Perera, 2010; Perera et al., 2011; Roberts and Petraglia 2015) that some part of the banana plant could have been used for ritual purposes, as happens ethnographically today, but additional investigations on this particular issue are required to support such a suggestion.

### 5.3.2. Wild rice

Comparison with multiple rice phytolith reference samples suggests strongly that Rock Shelter rice phytoliths were derived from *Oryza nivara* and/or *O. rufipogon* (SOM 2), but the criteria used here cannot fully separate the perennial *rufipogon* from the annual *nivara* (Fig. 9). Understanding the ecology of wild rice provides an opportunity to explore the relationship between human activity and the presence of wild rice in the archaeological context. The ecology of the wild rice species clearly indicates differing modes of wild rice exploitation for food since the prehistoric in South Asia (Munasinghe, 2007; Vaughan et al., 2008; Fuller and Qin, 2009; Fuller, 2011). *O. nivara* commonly grows in dry habitats and has large-scale seed production and so could have been easily gathered by prehistoric people. On the other hand, *O. rufipogon* prominently grows in aquatic habitats in relatively small isolated patches, not in genetically uniform stands and has a much lower seed production, so would have been more difficult to gather in quantity (Claridge et al. 1982; Vaughan, 1994; Vaughan and Morishima, 2003; Vaughan et al., 2008; Fuller and Qin, 2009).

In Marine Isotope stage 3 (MIS3), the elevated  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  from Batadomba-lena rock shelter faunal remains (Roberts et al., 2017) and Himalayan ice cores (Thompson et al., 2000) indicate that during the period 36,000-29,000 cal. BP the lowland rainforests of Southern Asia were more open, with decreased canopy cover as a result of decreased rainfall and temperature (cf. Mercader, 2002). Then, prior to the Last Glacial maximum (LGM), humid environments appear to have prevailed in the Indian subcontinent (Rajagopalan et al., 1997; Patnaik et al., 2008; Kumaran et al., 2013). Paleoclimatic records suggest that atmospheric cooling by 3-4°C occurred in the Tropics (Farrera et al., 1999), with a remarkable drop in precipitation during the LGM much greater than during any of the earlier (e.g. middle Pleistocene) glaciations (Robert et al., 2017). Palaeoclimate data from Sri Lanka suggest a drier LGM punctuated by short sharp climatic ameliorations (Premathilake, 2012ab, 2015a). It is probable that climatic fluctuations within prevailing dry conditions during the late Quaternary may have resulted in a number of climatically adapted wild rice populations (cf. Premathilake, 2003; Premathilake and Risberg, 2003; Fuller, 2011). Gathering by prehistoric humans could have modified *O. nivara* populations leading to a more reliable wild grain source for human use, especially when they were already widely growing in ideal habitats associated with

prolonged dry conditions long before the domesticated forms arose (Fuller et al., 2007; Fuller and Qin, 2009, Fuller et al., 2009; Fuller et al., 2010). During early rice exploitation, it is worth noting that high microcharcoal, and pollen and phytolith core records indicate regular anthropogenic burning from the Terminal Pleistocene (14,500-13,000 cal. BP) through the early-middle Holocene in the archaeological sites from Ganges plains, India (Singh, 2005; Saxena et al., 2006). Several archaeological sites in the Yangtze valley in China, dating from 17,000 cal. BP through the Terminal Pleistocene yielded *O. nivara* phytoliths (Zhijun, 1998; Chi, 2002; Lu, 2006; Liu et al., 2007; Zong et al., 2007). Sediment and peat core records from the Horton Plains, central Sri Lanka yielded *Oryza* spp. phytoliths and other evidence including pollen, charcoal, mineral magnetism, stable carbon and diatoms suggesting that burning and disturbance were associated with dry climate in montane environments occurred between 15,900-13,800 cal. BP (Premathilake, 2003, 2006; 2015a; Premathilake and Gunathilaka, 2013). Those records indicate that wild rice was present in human economies from the late Pleistocene into the Holocene in South and East Asia. We argue that wild rice exploited by late Pleistocene hunter-gatherers in Sri Lanka was *O. nivara* (rather than *O. rufipogon*) as an adaptation to the predominantly dry climate. The antiquity of human use of wild rice species, probably *O. nivara* at Fahien is remarkably early at 44,952-47,854 cal. BP, compared with the evidence for rice foraging in other known South Asian sites (Fuller, 2006; Premathilake, 2015b; Fuller et al., 2007).

### 5.3.3. Wild rice (*Oryza* spp.)

Leaf and seed (glum) phytolith evidence of wild rice, together with low counts of chlorodoid (e.g. *Cynodon* sp.), panicoid (e.g. *Panicum* sp.), and sedge (Cyperaceae), taxa which are recurrent weeds in dry rice fields with *O. nivara* (Table S1, S3; Weisskopf, et al., 2014; Huan et al., 2015; Jenkins et al., 2017) suggest that dry wild rice stands located near to the Rock Shelter were exploited. The samples analyzed yielded low numbers of phytoliths of dicotyledons and relatively high numbers of burnt phytoliths. These Rock Shelter phytolith records appear to be indicative of the conditions associated with anthropogenic activity, perhaps removal of competing vegetation and burning (cf. Jenkins et al., 2017).

The common husk phytoliths (single and double peaked forms, glumes, dendriforms) make it highly likely that wild rice seeds were exploited for food. The occurrence of burnt bulliforms and bilobates from the rice leaves and phytoliths from rice husks in association with ash deposits throughout the sequence (Table 1; Table S1, S3) suggests that rice straw was imported into the Rock Shelter. The presence of some bulliform type B, elongate psilate, elongate echinate, pooid, chloridoid and panicoid phytoliths in these assemblages indicates that wild grasses were also brought into the Rock Shelter. It is possible that wild rice and contaminant grassy plants were imported for threshing, but these may alternatively have been gathered for bedding, flooring, mat-making, thatching, ritual or other purposes (Tsartsidou et al., 2007; Pearsall, 2015; Bates et al., 2017; Premathilake et al., 2017a; Jenkins et al., 2017). From these primary uses, some may have ended up as fuel for domestic fires or may have been burnt to discourage insect pests, analogous with *fumier* deposits in Mediterranean caves (Angelucci et al. 2009).

The Rock Shelter rice phytolith assemblages thus may reflect human activities related to the model of prehistoric rice exploitation (e.g. Harvey and Fuller, 2005; Fuller and Qin, 2009; Fuller et al., 2011).

#### 5.3.4. Wild breadfruit (*Artocarpus nobilis*)

Samples from the Rock Shelter sediments contained phytoliths of wild breadfruit. In modern times, breadfruit is an important economic crop in South and Southeast Asia. (Dassanayake and Fosberg, 1981; Jarrett, 1959; Blench, 2004). Wild breadfruit, *Artocarpus nobilis*. is endemic to Sri Lanka, occurring in disturbed lowland rainforest from 600 to 800 m asl (Bandaranaike and Sultanbawa, 1991). Seeded wild breadfruit is an excellent source of carbohydrates and also contains significant amounts of geranylated phenolic compounds which have strong antioxidant properties (Jayasinghe et al., 2006). The seeds contain an especially high percentage of oil, which is commonly used in traditional medical practices. Since historic times seeded fruits have been one of the staple starchy foods in the traditional diet, providing a secure food source (Rajapaksha, 1998). The processing and preservation techniques for wild breadfruit include roasting, boiling and drying. The wood of the plant is also commonly used for fuel and for building material.



Charred tissues from fruits and seeds of wild breadfruit have been found from 36,000 cal. BP through to the terminal Pleistocene in the Batadomba-lena and Beli-lena rock shelters (Kajale, 1989; Deraniyagala, 1992; Perera et al., 2011). At Fahien, phytoliths suggest that the late Pleistocene rainforest hunter gatherers exploited seedy fruits of wild breadfruit as early as 44,952-47,854 cal. BP. (Fig. 10). Burnt phytoliths associated with charcoal and ash deposits recovered from the Rock Shelter clearly indicate that material from *Artocarpus* cf. *nobilis* was probably used as fuel (Perera, 2010; Table 1; Table S1, S3). We contrast the phytoliths of wild breadfruit at Fahien, with the morphologically different forms from two other breadfruit species, *Artocarpus altilis* (rata del), which was probably domesticated in New Guinea (Blench, 2004) and *Artocarpus heterophyllus* (kos), native to Western part of India (Rajapaksha, 1998). Both species are commonly cultivated in Sri Lanka today.

#### 5.3.5. *Durian* (*Durio zibethius*)

This is a lowland rainforest species, usually utilized as desert fruits, very delicious to some people, although the wood is also valued for construction. It is widely cultivated throughout Sri Lanka, Indonesia, Malaysia, India and Burma, and contains significant amounts of carbohydrates, proteins, fats, calcium, phosphorus, vitamins (A, B1, B2, C and E) and fibre (Rajapaksha, 1998). *Durio zibethius* perhaps originated in Borneo (Blench, 2004, 2008), and spread throughout the Malayan peninsula during the pre-European period and then to a number of other countries in South and Southeast Asia during the nineteenth century. It is interesting to note that no phytolith and other archaeobotanical evidence of *Durio* spp. from any cave and rock shelter excavation sites in South and Southeast Asia has been reported (cf. Blench, 2004, 2008). Phytolith evidence for *Durio* cf. *zibethius* was identified in later Pleistocene samples (Fig. 10) and pollen from *Durio* sp. was also found in all the samples analyzed (to be published elsewhere). However, phytolith evidence of *Durio* cf. *zibethius* at Fahien Rock Shelter is somewhat striking compared to the available records (Blench, 2004, 2008), requiring additional investigations on a number of well-dated sites to gain a better understanding the archaeobotany of *Durio* cf. *zibethius* in the rainforests of South and Southeast Asia. Nevertheless, evidence for this economically-important plant is consistent with finds of phytoliths of other food plants.

#### 5.3.6. *Canarium* sp. nuts

Phytoliths classified as from the Burseraceae are closely consistent with the modern phytoliths and macro archaeobotanical remains of *Canarium* sp. nuts suggesting that this was part of the subsistence pattern of the Rock Shelter occupants (Table 1; Table S1, S3; Fig. 7; Fig. 10). *Canarium zeylanicum* (Burseraceae family), an endemic tree species commonly grows in the lowland rainforest of Sri Lanka (Gunatilleke et al., 2008). Seeds of the species found in Batadomba-lena rock shelter (Deraniyagala 1992; Perera 2010; Perera et al. 2011) and Belilena (Kajale, 1989; Deraniyagala 1992) were dated to 16,000 cal. BP and the terminal Pleistocene age respectively. Records of *Canarium* sp. dated through late Pleistocene to early Holocene (14,000-10,000 cal. BP) from Sepik-Ramu in New Guinea (Latinis, 2000), Con Moong cave in Vietnam (Viet, 2001), Spirit cave in Thailand (Gorman, 1970) and the older record ~45,000 cal. BP from Niah Cave in Sarawak (Paz, 2005) suggest that *Canarium* spp. was integral to subsistence Near Oceania, South and Island Southeast Asia. In an environmental context, the appearance of Burseraceae phytoliths at Fahien and an increasing occurrence of *Canarium* sp. nuts at other Sri Lankan sites indicates an expansion of the lowland rainforest (Roberts et al., 2017), which corresponds to the strengthening of southwest monsoon rains through the terminal Pleistocene to the early Holocene (cf. Premathilake, 2003, 2006, 2012ab; 2015a; Premathilake and Gunathilaka, 2013; Perera et al., 2011).

#### 5.3.7. *Species of Palmae*

An abundance of Palmae phytoliths, identified as Palmae 1 and 2 based on morphological variations, indicates that the Rock Shelter occupants brought the vegetative parts of palms into the Rock Shelter. Palmae 1 are spheroid-elliptical (around 10-30  $\mu$ m), densely echinate phytoliths while Palmae 2 includes spheroid-elliptical (5-10  $\mu$ m) phytoliths with nodular obtuse tip projections (Fig. 10). In our reference collection, phytoliths of Palmae 1 are broadly similar to the phytoliths derived from *Borassus* sp., while Palmae 2 is similar to the phytoliths from *Areca* sp., *Caryota* sp., *Cocos* sp. and *Phoenix* sp. However, the morphological differences within these two groups are difficult to apply to species level identification. It is possible to suggest that blackened and non-blackened Palmae phytoliths in association with ash deposits throughout the sequence were released from palm leaves, bark

and stems used by Rock Shelter occupants, possibly for fire, food, vessels and construction as early as 44,952-47,854 cal. BP.

In modern times, several palm species are frequently associated with human activity (Henderson, 2009). Eighteen native species are found in Sri Lanka, ten of which are endemic and several native palms grow in the disturbed lowland rainforest, e.g. at Sinharaja Forest Reserve (Henderson, 2009). Ethno-archaeological studies indicate that palm leaves are traditionally used for baskets, fishing nets, containers, mats, sleeping platforms (Premathilake et al., 1994). In south and southeast Asia, *Borassus flabellifer* (Palmyra palm) is now commonly used for roofing, sugar, toddy (naturally alcoholic sap of the palm) and writing (using the dried leaves), but the origins and original uses of the species during prehistoric times are disputed (Blench, 2005). Three palms, *Areca catechu*, *Caryota urens* and *Cocos nucifera* are among the most common economic crops. The first two grow naturally in disturbed lowland rainforest in Sri Lanka, India, Malaysia, Myanmar and Nepal. The third commonly grows in tropical and subtropical coastlands. Its origins are much disputed (Blench, 2005), however, linguistic studies suggest that it was carried to Sri Lanka and India prior to the second century BC (Mahdi, 1998). Harris (1990) suggests that coconut originated in Malaysia and its general distribution pattern is a relic of the splitting-up of Gondwanaland, highlighting the complexity of its botanical origins. Blench (2004, 2008) reports *Areca catechu* was in use for chewing at 13,000 cal. BP in Malaysia, and *Cocos nucifera* and *Areca catechu* occurred as part of an early agricultural system in Papua New Guinea at 6,000-6,950 cal. BP (Denham, 2004; Fairbairn and Swadling, 2005). Two coastal sites located in south- and north-western Sri Lanka yielded pollen evidence of *Cocos nucifera* dating to 7,500 cal. BP and 3,400 cal. BP respectively (Manawadu, 2016; Dilrukshi, 2017), confirming the antiquity of the accounts of coconut (Mahdi, 1998). Phytolith and starch grains from archaeological contexts dated to 5,000 cal. BP suggest that sago palms were an important plant food prior to rice in south subtropical China (Yang et al., 2013), and the Fishtail Palm (*Caryota* sp.) has been mentioned by Ki Han during the Jin Dynasty (AD 290-307) (Ruddle et al., 1978). Archaeobotanical records from North Gujarat (India) suggest that the progressive weakening of the south west monsoon around 7,000 cal. BP, compelled human populations to adapt seminomadic pastoralism which involved the cultivation of fast-maturing crops, and the gathering of a number of wild plants

including *Phoenix sylvestris* (García-Granero et al., 2016). In Sri Lanka, ashy deposits at the Batadomba-lena Rock Shelter identified as hearths yielded round echinate Palmae phytoliths, suggesting the use of Palmae-based artefacts (e.g. matting, thatching and woven ware) after the LGM (Perera et al., 2011).

#### 5.3.8. *Bamboo*

Phytoliths from the Fahien sediments imply that bamboos were abundant in the lowland rainforest. It is likely that Rock Shelter occupants may have used them for tools, containers, construction, fuel and other purposes as evidenced by an abundance of burned and un-burned rondel and large saddle phytoliths (Fig. 10) identical to those of modern Bambusoideae (**SOM 2**) Bamboos are perennial evergreen plants found in diverse climates from cold mountains to hot tropical regions and are especially common in lowland disturbed rainforest in Sri Lanka. They are of great cultural and economic significance, and are used as materials for various types of construction, tool making, baskets, vessels, mats, food, medicine, textiles and wood throughout South, East and Southeast Asia (Roach, 2013). Bulmer (1975) reported the presence of bamboo in early- to middle-Holocene archaeological contexts at Yuku, Papua New Guinea. Archaeological excavations in Japan uncovered bamboo baskets dating to the Late Jomon Period 4000-3000 cal. BP (Piper et al., 2017)

#### 5.39. *Diatoms*

The freshwater diatoms may indicate human activities; for instance from faeces, having been incorporated from aquatic microflora in drinking water, or from external contamination of the Rock Shelter occupants who have visited freshwater habitats for fishing and/or bathing, or on plants or animals taken from freshwaters (cf. Perera, 2010). Alternatively, the diatoms might represent the construction of floors using mud from flood plains or rivers in the vicinity of the Rock Shelter, although if this is the case no other evidence for this practice survives. Infrequent visits to the brackish/marine environment for fishing and salt collection may account for the discontinuous record of marine diatoms and is consistent with the finds of shark teeth and vertebra and marine shell beads through the late Pleistocene sequence at the Rock Shelter (Perera, 2010).

## 6. Conclusions

Understanding the human history of the rainforests of Sri Lanka is poor because of the lack of preservation of organic materials and effect of taphonomic processes. The Late Pleistocene archaeological sequence dating back to 44,952-47,854 cal. BP at the Fahien Rock Shelter in lowland southwest Sri Lanka was an excellent opportunity to supplement our understanding using siliceous phytoliths. The phytolith evidence indicates that the Rock Shelter occupants exploited wild bananas (*Musa acuminata* and *Musa balbisiana*), wild rice (most possibly *Oryza nivara*), wild breadfruit (*Artocarpus nobilis*), durians (*Durio* cf. *zibethius*), *Canarium* sp. and a number of species of palm and bamboo. Most of these are commonly associated with disturbed lowland rainforests. These results are of great importance as the oldest evidence for late Pleistocene rainforest plant exploitation from South Asia. This is relatively younger, however, than the earliest evidence from the Niah Cave, Sarawak dating to 53,000 cal. BP (Barker, 2011; Barker and Farr, 2016; Hunt et al., 2016; Barker et al, 2017). An uncertainty left by the current work is whether the earliest deposits sampled here actually relate to the first modern anatomical human presence in Sri Lanka. We assume that further excavation of the very lowermost layers at the Rock Shelter is likely to produce additional evidence in due course, and it would be most interesting to investigate the use of rainforest plants by earlier colonists.

## Acknowledgments

The British Academy provided a Visiting Fellowship for RP at the School of Geography, Palaeoecology and Archaeology, Queens University Belfast, where COH was the UK applicant. Financial support received from National Research Council (NRC, grant-14-43) is appreciated. We thank Professors Keith Bennett, Paula Reimer, Jonathan Pilcher and Dr. Maarten Blaauw for supporting the project. We thank Mrs. Elizabeth Woodgyer and Mr. Martin Xanthos of the Herbarium, Royal Botanic Garden, Kew, for providing herbarium specimens. Directions and field facilities received from Dr. Nimal Perera, Director Excavation and Dr. Senerat Dissanayake, Director General, Department of Archaeological Survey and Dr. S. U. Deraniyagala, Former Director General, Department of Archaeological Survey, Sri Lanka are greatly appreciated. Dr. David Simpson, QUB and Mrs. Sewwandhi Gunasekara, University of Colombo School of Computing are thanked for their constructive comments. RP thanks Mr.

W.M.C. Oshan, Chief Excavation Supervisor to the Fahien Project at the Sri Lanka Department of Archaeology, Mr. Asoka Perera and Mr. Sampath Perera at the Postgraduate Institute of Archaeology, University of Kelaniya for field support; the Director, Professor Jagath Weerasinghe, and the Board of Management at the PGIAR, University of Kelaniya for leave to complete this project and Mr. Jim Bradley for personal support.

## References

- Albert, R. M., Lavi, O., Estroff, L. Weiner, S., Tsatskin, A., Ronen, A., Lev-Yadun, S., 1999. Mode of Occupation of Tabun Cave, Mt Carmel, Israel During the Mousterian Period: A Study of the Sediments and Phytoliths. *Journal of Archaeological Science* 26 (10), 1249-1260.
- Albert, R.M., Bamford, M.K., Cabanes, D., 2006. Taphonomy of phytoliths and macroplants in different soils from Olduvai Gorge (Tanzania) and the application to Plio-Pleistocene paleoanthropological samples. *Quaternary International* 148, 78-94.
- Alexandre, A., Meunier, J. D., Colin, F., Koud, J. M., 1997. Plant impact on the biogeochemical cycle of silicon and related weathering processes. *Geochimica et Cosmochimica Acta* 61, 677-682.
- Alexandre, A., Basile-Doelsch, I., Delhay, T., Borshneck, D., Mazur, J.C., Reyerson, P., Santos, G.M., 2015. New highlights of phytolith structure and occluded carbon location: 3-D X-ray microscopy and NanoSIMS results. *Biogeosciences*, 12, 863-873.
- Angelucci, D.E., Boschian, G., Fontanals, M., Pedrotti, A., Vergès, J.M. 2009. Shepherds and karst: the use of caves and rock-shelters in the Mediterranean region during the Neolithic, *World Archaeology*, 41, 2, 191-214.
- Bandaranaike, W. M., Sultanabawa, M.U.S., 1991. A list of Endemic Plants of Sri Lanka. Forest Department, Colombo, Sri Lanka.
- Barker G. 2011. The cost of cultivation. *Nature* 473: 164-164.
- Barker G, Farr L. 2016. Archaeological investigations in the Niah Cave, Sarawak. (McDonald Institute for Archaeological Research, University of Cambridge, Downing Street, Cambridge.
- Barker, G., Barton, H., Bird, M., Daly, P., Datan, I., Dykes, A., Farr, L., Gilbertson, D., Harisson, B., Hunt, C., Higham, T., Krigbaum, J., Lewis, H., McLaren, S., Paz, V., Pike, P., Piper, P., Pyatt, B., Rabett, R., Reynolds, T., Rose, J., Rushworth, G., Stephens, M., Stringer, C., Thompson, G., 2007. The human revolution in tropical Southeast Asia: the antiquity of anatomically modern humans, and of behavioural modernity, at Niah Cave (Sarawak, Borneo). *Journal of Human Evolution* 52, 243-261.
- Barker, G., Hunt, C., Barton, H., Gosden, C., Jones, S., Lloyd-Smith, L., Farr, L., Nyirí, B., O'Donnell, S., 2017. The 'cultured rainforests' of Borneo. *Quaternary International* 448, 44-61.
- Bates, J., Singh, R. N., Petrie, C. A., 2017. Exploring Indus crop processing: combining phytolith and macrobotanical analyses to consider the organization of agriculture in northwest India c. 3200-1500 BC. *Vegetation History Archaeobotany* 26, 25-41. DOI 10.1007/s00334-016-0576-9.
- Behrensmeier, A. K., Kidwell, S. M., Gastaldo, R. A., 2000. Taphonomy and Paleobiology. *Paleobiology* 26, 4, Supplement 103-144.

- Blench, R., 2004. Fruits and arboriculture in the indo-pacific region. Indo-pacific prehistory association bulletin 24, 31-50.
- Blench, R., 2005. Fruits and arboriculture in the Indo-Pacific region. Bulletin of the Indo-Pacific Prehistory Association 24, 31-50.
- Blench, R., 2008. A history of fruits on the Southeast Asian mainland. In: Toshiki, O., Akinori, U. (Eds), Occasional Paper 4 Linguistics, Archaeology and the Human Past, Research Institute for Humanity and Nature, Kyoto, Japan, pp.115-135.
- Borrelli, N., Alvarez, M.F., Osterrieth, M.L., Marcovecchio, J.E., 2010. Silica content in soil solution and its relation with phytolith weathering and silica biogeochemical cycle in Typical Argiudolls of the Pampean Plain, Argentina; a preliminary study. Journal of Soils Sediments 10, 983-994.
- Bowdery, D., 1999. Phytoliths from Tropical Sediments: Reports from Southeast Asia and Papua New Guinea. Indo-Pacific Prehistory Association Bulletin 18, 159-168.
- Bulmer, S., 1975. Settlement and Economy in Prehistoric Papua New Guinea: a Review of the Archaeological Evidence. Journal de la Société des Océanistes, 31, 7-75.
- Butzer, K. W., 2008. Challenges for a cross-disciplinary geoarchaeology: The intersection between environmental history and geomorphology. Geomorphology 101, 402-411.
- Cabanes, D., Weiner, S., Shahack-Gross, R., 2011. Stability of phytoliths in the archaeological record: a dissolution study of modern and fossil phytoliths. Journal of Archaeological Science 38, 2480-2490.
- Cabanes, D., Gadot, Y., Cabanes, M., Finkelstein, I., Weiner, S., Shahack-Gross, R., 2012. Human impact around settlement sites: a phytolith and mineralogical study for assessing site boundaries, phytolith preservation, and implications for spatial reconstructions using plant remains. Journal of Archaeological Science 39, 2697-2705.
- Castillo, C., Fuller, D. Q., 2010. Still too fragmentary and dependent upon chance? Advances in the study of early Southeast Asian archaeobotany. In: Bellina-Pryce, B., Pryce, T.O., Bacus, E., Wisseman-Christie, J. (Eds.), 50 Years of Archaeology in Southeast Asia: Essays in Honour of Ian Glover, Bangkok, River Books, pp. 90-111.
- Castillo, C., Fuller, D.Q., 2015. Bananas: The Spread of a Tropical Forest Fruit as an Agricultural Staple. In: Lee-Thorp, J., Katzenberg, M.A. (Eds.), Oxford Handbooks of the Archaeology of Diet, Oxford University Press, pp. 1-26.
- Clarkson, C., Petraglia, M., Korisettar, R., Haslam, M., Boivin, N., Crowther, A., Ditchfield, P., Fuller, D.Q., Miracle, P., Harris, C., Connell, K., James, H., Koshy, J., 2009. The oldest and longest enduring microlithic sequence in India: 35,000 years of modern human occupation and change at the Jwalapuram Locality 9 rock shelter. Antiquity 83, 326-348.
- Chandrarathna, M. F., Nanayakkara, K.D.S.S., 1951 .Cultivated varieties of banana in Ceylon. Tropical Agriculturist 107, 70-91.
- Cheeseman, E.E. 1947. The classification of the bananas. Kew Bulletin 2, 106-117.
- Chi, Z., 2002. Early pottery and rice phytoliths remains from Xianrendong and Diaotonghuan sites, Wannian, Jiangxi Province. In: Yasuda, Y. (Ed.), Pottery and Agriculture, Roli Book Pvt. Ltd., India, pp. 185-191.
- Christensen, H., 2002. Ethnobotany of the Iban & the Kelabit. Forest Department Sarawak, Malaysia; NEPCon; University of Aarhus, Kuching and Aarhus.
- Claridge, M.F., Denhollander, J., Furet, I., 1982. Adaptation of brown plantho (*Nilaparvata lugens*) populations to rice varieties in Sri Lanka. Entomologia Experimentalis et Applicata 32, 222-226. DOI: 10.1111/j.1570-7458.1982.tb03209.x

- 714 Cooray, P.G., 1984. An introduction to the Geology of Sri Lanka. National Museum Publications,  
715 Colombo.
- 716 Cosgrove, R., Field, J., Ferrier, Å. 2007. The archaeology of Australia's tropical rainforests.  
717 Palaeogeography, Palaeoclimatology, Palaeoecology 251, 150-173
- 718 Constantine, D., 1999-2008. The Musaceae: An annotated list of the species of *Ensete*, *Musa* and  
719 *Musella*. [www.users.globalnet.co.uk/~drc/musaceae.htm](http://www.users.globalnet.co.uk/~drc/musaceae.htm). (Last accessed 20.1.2009).
- 720 Dassanayake, M.D., Fosberg F.R., 1981. A revised handbook to the flora of Ceylon 3, Oxford  
721 and IBH, New Delhi.
- 722 De Langhe, E., 2009. Relevance of banana seeds in archaeology. *Ethnobotany Research &*  
723 *Applications* 7, 271-281.
- 724 Denham, T. P., Haberle, S.G., Lentfer, C., Fullagar, R., Field, J., Therin, M., Porch, N.,  
725 Winsborough, B., 2003. Origins of agriculture at Kuk Swamp in the Highlands of New  
726 Guinea. *Science* 301, 189-193.
- 727 Denham T. 2004. The roots of agriculture and arboriculture in New Guinea: looking beyond  
728 Austronesian expansion, Neolithic packages and indigenous origins. *World Archaeology* 36,  
729 610-620.
- 730 Deraniyagala, S. U., 1992. The Prehistory of Sri Lanka: An Ecological Perspective, second ed.  
731 Department of Archaeological Survey, Colombo.
- 732 Dilrukshi, W.D.J., 2017. Palaeoecological events stratigraphies recorded by organic carbon and  
733 palynological sequences from Sirigampola, southwest Sri Lanka. Postgraduate Institute of  
734 Archaeology, University of Kelaniya, Sri Lanka (Master of Philosophy Thesis)
- 735 Fairbairn, A., Swadling, P., 2005. Re-dating mid-Holocene betelnut (*Ereca catechu*) and other  
736 plant use at Dongan, Papua New Guinea. *Radiocarbon* 47, 3, 377-382.
- 737 Farmer, V. C., Delbos, E., Miller, J. D., 2005. The role of phytolith formation and dissolution in  
738 controlling concentrations of silica in soil solutions and streams. *Geoderma* 127, 71-79.
- 739 Farrera, I., Harrison, S.P. et al. 1999. Tropical climates at the Last Glacial Maximum: a new  
740 synthesis of terrestrial palaeoclimate data, vegetation, lake levels and geochemistry. *Climate*  
741 *Dynamics* 15, 823-856.
- 742 Fraysse, F., Cantais, F., Pokrovsky, O.S., Schott, J., Meunier, J.D., 2006. Aqueous reactivity of  
743 phytoliths and plant litter: physico-chemical constraints on terrestrial biogeochemical cycle of  
744 silicon. *Journal of Geochemical Exploration*. 88, 202-205.
- 745 Fraysse, F., Pokrovsky, O.S., Schott, J., Meunier, J.-D., 2009. Surface chemistry and reactivity of  
746 plant phytoliths in aqueous solutions. *Chemical Geology* 258, 197-206.
- 747 Fredlund, G. G., Tieszen, L.L., 1997. Phytolith and carbon isotope evidence for late quaternary  
748 vegetation and climate change in the southern Black Hills, South Dakota. *Quaternary*  
749 *Research* 47, 206-217.
- 750 Friesem, D. E., Lavi, N., Madella, M., Ajithprasad, P., French, C., 2016. Site Formation  
751 Processes and Hunter Gatherers Use of Space in a Tropical Environment: A Geo-  
752 Ethnoarchaeological Approach from South India. *PLoS ONE* 11, 10, 1-30. PLOS ONE |  
753 DOI:10.1371/journal.pone.0164185.
- 754 Fuller, D. Q., 2006. Agricultural origins and frontiers in South Asia: a working synthesis. *Journal*  
755 *of World Prehistory*, 20, 1-86.
- 756 Fuller, D. Q., Harvey, E. L., 2006. The archaeobotany of Indian pulses: identification, processing  
757 and evidence for cultivation. *Environmental Archaeology* 2, 219-246.



758 Fuller, D. Q, Harvey, E. L, Qin, L. 2007. Presumed domestication? Evidence for wild rice  
759 cultivation and domestication in the fifth millennium BC of the Lower Yangtze region.  
760 *Antiquity*, 81, 316-31.

761 Fuller, D. Q., Madella, M., 2009. Banana Cultivation in South Asia and East Asia: A review of  
762 the evidence from archaeology and linguistics. *Ethnobotany Research and Applications* 7,  
763 333-351.

764 Fuller, D.Q., Qin, L., Zheng, Y., Zhao, Z., Chen, X., Hosoya, L.A., Sun, G., 2009. The  
765 domestication process and domestication rate in rice: Spikelet bases from the Lower Yangtze.  
766 *Science* 323, 1607-1610.

767 Fuller, D.Q., Qin, L., 2009. Water management and labour in the origins and dispersal of Asian  
768 rice. *World Archaeology* 41, 1, 88-111.

769 Fuller, D.Q., Sato Y-I, Castillo, C., Qin, L., Weisskopf, A.R., Kingwell-Banham, E.J., Song, J.,  
770 2010. Consilience of genetics and archaeobotany in the entangled history of rice.  
771 *Archaeological and Anthropological Sciences* 2(2), 115-131.

772 Fuller, D.Q., Etten, J. V., Manning, K., Castillo, C, Kingwell-Banham, E., Weisskopf, A., Qin, L.,  
773 Sato, Y., Hijmans, R. J., 2011. The contribution of rice agriculture and livestock pastoralism to  
774 prehistoric methane levels: An archaeological assessment. *The Holocene* 21 (5), 743-759.

775 Fuller, D.Q. 2011. Pathways to Asian Civilizations: Tracing the Origins and Spread of Rice and  
776 Rice Cultures. *Rice* 4, 78-92. DOI 10.1007/s12284-011-9078-7.

777 García-Granero, J. J., Lancelotti, C., Madella, M., Ajithprasad, P., 2016. Millets and Herders:  
778 The Origins of Plant Cultivation in Semiarid North Gujarat (India). *Current Anthropology* 57,  
779 2, 149-173.

780 Gérard, F., Mayer, K. U., Hodson, M. J., Ranger, J., 2008. Modeling the biogeochemical cycle of  
781 silicon in soils: application to a temperate forest ecosystem. *Geochimica et Cosmochimica*  
782 *Acta* 72, 741-758.

783 Gilbertson, D., Bird, M., Hunt, C., McLaren, S., Mani Banda, R., Pyatt, B., Rose, J. & Stephens,  
784 M. 2005. Past Human activity and geomorphological change in a guano-rich tropical cave  
785 mouth: initial interpretations of the Late Quaternary succession in the Great Cave of Niah,  
786 Sarawak. *Asian Perspectives*, 44, 1, 16-41.

787 Goldberg, P., Bar-Yosef, O., 1998. Site formation processes in Kebara and Hayonim Caves and  
788 their significance in Levantine prehistoric caves. In: Akazawa, T., Aoki, K., Bar-Yosef, O.  
789 (Eds.), *Neanderthals and Modern Humans in Western Asia*. New York: Plenum Press, pp.  
790 107-26.

791 Gorman, C.F., 1970. Excavations at Spirit Cave, North Thailand: Some Interim Interpretations.  
792 *Asian Perspectives* 13, 79-107.

793 Gunatilleke, N., Pethiyagoda, R., Gunatilleke, S., 2008. Biodiversity of Sri Lanka. *Journal of*  
794 *National Science Foundation Sri Lanka* 36, 25-62

795 Gunatillake, C.V. S, Gunatillake, A. U. N., Ethugala A. U. K., Esufali, S. 2004. Ecology of  
796 Sinharaja Rain Forest and the Forest Dynamics Plot in Sri Lanka's Natural World Heritage  
797 Site. WHT Publications (Pvt.) Ltd.

798 Gunatilaka A. 2007. Role of basin-wide landslides in the formation of extensive alluvial  
799 gemstones in Sri Lanka. *Earth Surface Processes and Landforms*, 32, 1863–1873.

800 Häkkinen, M., Väre, H., 2008. Typification and check-list of *Musa* L. names (Musaceae) with  
801 nomenclatural notes. *Adansonia série* 3, 30, 63-112.

802 Harries, H. C., 1990. Malaysian origin for a domestic *Cocos nucifera*. In: Baas, P., Kalkman, K.,  
803 Geesink, R. (Eds.), *The Plant Diversity of Malaysia*. Kluwer, Dordrecht, pp.351-357.

- Harvey, E.L., Fuller, D. Q. 2005. Investigating crop processing using phytolith analysis: the example of rice and millets. *Journal of Archaeological Science* 32, 739-752.
- Haslam, M., Clarkson, C., Roberts, R.G., Bora, J., Korisettar, R., Ditchfield, P., Chivas, A.R., Harris, C., Smith, V., Oh, A., Eksambekar, S., Boivin, N., Petraglia, M., 2012. A southern Indian Middle Palaeolithic occupation surface sealed by the 74 ka Toba eruption: further evidence from Jwalapuram Locality 22. *Quaternary International* 258, 148-164.
- Henderson, A., 2009. *Palms of Southern Asia*. Princeton University Press, USA.
- Heyne, K., 1950. *De Nuttige Planten van Indonesie*. 3<sup>rd</sup> Ed. Jakarta, Indonesia, Wageningen.
- Headland, T. N., Bailey, R. C., 1991. Introduction: Have Hunter-Gatherers Ever Lived in Tropical Rain Forest Independently of Agriculture? *Human Ecology*, 19, 2, 115-122.
- Huan, X., Lu, H., Wang, C., Tang, X., Zuo, X., Ge, Y., He, K., 2015. Bulliform Phytolith Research in Wild and Domesticated Rice Paddy Soil in South China. *PLoS ONE* 10(10): e0141255. doi:10.1371/journal.pone.0141255
- Hunt, C., Davison, J., Inglis, R., Farr, L., Barker, G., Reynolds, T., Simpson, D., el-Rishi, H. 2010. Site formation processes in caves: the Holocene sediments of the Haua Fteah, Cyrenaica, Libya. *Journal of Archaeological Science* 37, 1600–1611.
- Hunt, C. O., Gilbertson, D.D., Rushworth, G., 2012. A 50,000-year record of late Pleistocene tropical vegetation and human impact in lowland Borneo. *Quaternary Science Reviews* 37, 61-80.
- Hunt, C., Kealhofer, L., Premathilake, R., Garry, R., Gilbertson, D., Jones, S., Thompson, G., 2016. Palynology, phytoliths, diatoms and wood in the west mouth: stratigraphic and taphonomic studies of late Quaternary vegetation history. In: Barker, G., Farr L., (Eds.), *Archaeological investigations in the Niah Caves, Sarawak*, pp. 149-176.
- Jarrett, F.M., 1959. Studies on *Artocarpus* and allied genera. A revision of subgenus *Artocarpus*. *Journal of Arnold arboretum* 60, 327-368.
- Jayasinghe, J., Rupasinghe, G. K., Hara, N., Fujimoto, Y., 2006. Geranylated phenolic constituents from the fruits of *Artocarpus nobilis*. *Phytochemistry* 67, 13, 1353-1358.
- Jenkins, E. L., Allcock, S. L., Elliott, S., Palmer, C., Grattan, J., 2017. Ethno-geochemical and phytolith studies of activity related patterns: a case study from Al Ma'tan, Jordan. *Environmental Archaeology* 22, 4, 412-433.
- Karkanas, P., 2010. Preservation of anthropogenic materials under different geochemical processes: a mineralogical approach. *Quaternary International* 214, 63-69.
- Kajale, M. D., 1989. Mesolithic exploitation of wild plants in Sri Lanka: archaeobotanical study at the cave of Beli-Lena. In: Harris, D.R., Hillman, G. C. (Eds.), *Foraging and Farming: The Evolution of Plant Exploitation*. Unwin Hyman, London, pp. 269-281.
- Kennedy, J., 2009. Bananas and People in the Homeland of Genus *Musa*: Not just pretty fruit. *Ethnobotany Research & Applications* 7, 179-197.
- Kinnaird, T. C., Sanderson, D. C. W., 2010. Luminescence Dating of Sediments from the Fahien-lena Rock shelter, Southern Sri Lanka. Technical Report, Scottish Universities Environmental Research Centre, East Kilbride, UK.
- Kourampas, N., Simpson, I. A., Perera, N., Deraniyagala, S. U., Wijeyapala, W. H., 2009. Rock Shelter sedimentation in a dynamic tropical landscape: late Pleistocene to Early Holocene archaeological deposits in Kutulgala Beli-lena, southwestern Sri Lanka. *Geoarchaeology* 24, 677-714.

- Kumaran, K.P.N., Limaye, R.B., Punekar, S.A., Rajaguru, S.N., Joshi, S.V., Karlekar, S.N., 2013. Vegetation response to South Asian monsoon variations in Konkan, western India during the late Quaternary: evidence from fluvio-lacustrine archives. *Quaternary International* 286, 3-18.
- Latinis, D. K., 2000. The development of subsistence system models for Island Southeast Asia and Near Oceania: the nature and role of arboriculture and arboreal-based economies. *World Archaeology* 32,1, 41-67.
- Lentfer, C. J., Boyd, W. E., 1998. A comparison of three methods for the extraction of phytolith from sediments. *Journal of Archaeological Science* 25, 1159-1183.
- Lentfer, C., Pavlides, C., Specht, J., 2010. Natural and human impacts in a 35 000-yr vegetation history in central New Britain, Papua New Guinea. *Quaternary Science Reviews* 29, 3750-3767.
- Liu, L., Lee, G.A., Jiang, L., Zhang, J., 2007. Evidence for the early beginning (c. 9000 cal. BP) of rice domestication in China: a response. *The Holocene*, 17, 1059-1068.
- Loucaides, S., Van Cappellen, P., Behrends, T., 2008. Dissolution of biogenic silica from land to ocean: role of salinity and pH. *Limnology and Oceanography* 53, 1614-1621.
- Loucaides, S., Behrends, T., Van Cappellen, P., 2010. Reactivity of biogenic silica: surface versus bulk charge density. *Geochimica et Cosmochimica Acta* 74, 517-530.
- Lu, T. L. D. 2006. The occurrence of cereal cultivation in China. *Asian Perspectives*, 45: 129-58.
- Madella, M., Lancelotti, C., 2012. Taphonomy and phytoliths: A user manual. *Quaternary International* 275, 76-83
- Mahdi, W., 1998. Linguistic data on transmission of Southeast Asian cultigens to India and Sri Lanka. In: Blench, R.M., Spriggs, M. (Eds.), *Archaeology and Language II*. Routledge, London, pp.390-415.
- Manawadu, B., 2016. Middle-late Holocene environment, climate and human activities as recorded by palyno-stratigraphy from Villu ecosystem in Wilpattu National park, Northwestern Sri Lanka., Postgraduate Institute of Archaeology, University of Kelaniya, Sri Lanka. (Master of Science Thesis).
- Mercader, J. (Ed.), 2002. *Under the Canopy: The Archaeology of Tropical Rain forests*. Rutgers University Press, London.
- Munasinghe, M. L. A. M. S., 2007. Pollen morphological studies of selected cultivated and wild species of Poaceae. Postgraduate Institute of Archaeology, University of Kelaniya, Sri Lanka. (Master of Philosophy Thesis).
- Oshan, W. M. C., 2011. Stratigraphical analysis of Fahien Rockshelter in Sri Lanka: comprising stratigraphy excavated in 1988 and 2009. Department of Archaeology, Deccan College, Postgraduate Research Institute, Pune, India (Master of Art Dissertation).
- Osterrieth, M., Madella, M., Zurro, D., Alvarez, M. F., 2009. Taphonomical aspects of silica phytoliths in the loess sediments of the Argentinean Pampas. *Quaternary International* 193, 707-779.
- Patnaik, R., Badam, G. L., Murty, M. L. K., 2008. Additional vertebrate remains from one of the late Pleistocene–Holocene Kurnool caves (Muchchatla Chintamanu Gavi) of South India. *Quaternary International* 192, 43-51.
- Paz, V. J., 2005. Rock Shelters, Caves, and Archaeobotany in Island Southeast Asia, *Asian Perspectives* 44, 107-118.
- Pearsall, D. M., 2015. *MU Phytolith Classification System: Notes and Guidelines for Use*. Department of Anthropology, University of Missouri, Columbia.

- Perera, H. N., 2010. Prehistoric Sri Lanka: Late Pleistocene Rockshelters and an Open Air Site. BAR International Series. Archaeopress, Oxford.
- Perera, N., Kourampas, N., Simpson, I.A., Deraniyagala, S.U., Bulbeck, D., Kamminga, J., Perera, J., Fuller, D.Q., Szabó, K., Oliveira, N.V., 2011. People of the ancient rainforest: Late Pleistocene foragers at the Batadomba-lena rockshelter, Sri Lanka. *Journal of Human Evolution* 61, 254-269.
- Perera, H. N., 2015. The Importance of Sri Lanka Wet Zone Rockshelters. In: Dissanayake, S., Rev. Chanaloka, P., Kodituwakku, N., (Eds.), Piyasatahan, Department of Archaeological Survey and Ministry of Cultural Affairs, Colombo, pp.104-117.
- Perera, P., 2017. A record of modern phytoliths morphology from *Musa* bananas: an approach to identify banana phytoliths on archaeological context in Sri Lanka. Postgraduate Institute of Archaeology, University of Kelaniya, Sri Lanka. (Master of Philosophy Thesis)
- Petraglia, M., Clarkson, C., Boivin, N., Haslam M., Korisettar, R., Chaubey, G., Ditchfield, P., Fuller, D., James, H., Jones S., Kivisild, T., Koshy, J., Mirazon, Lahr, M.M., Metspalu, M., Roberts, R., Arnold, L., 2009. Population increase and environmental deterioration correspond with microlithic innovations in South Asia ca. 35,000 years ago. *Proceedings of the National Academy of Sciences* 106, 1226-12266.
- Piperno, D. R., 1988. Phytolith analysis: An archaeological and geological perspective. San Diego: Academic Press.
- Piperno, D. R., 2006. Phytoliths: a comprehensive guide for archaeologist and paleoecologist. Alta Mira press, Rowman and Littlefield Publishers, Inc. UK.
- Piper, P. J., Matsumura, H., Bulbeck, D., 2017. New perspectives in Southeast Asian and Pacific prehistory. The Australian National University, Acton ACT 2601, Australia.
- Premathilake, R., Epitawatta, D.S., Yasapala, N., Runebjer, E.M., 1994. Plants used in household. In: Bandaranayake, S., Epitawatta, S., Morgren, M. (Eds.), Further Studies in the Settlement Archaeology, PGIAR, Sri Lanka, pp. 219-232.
- Premathilake, R. 2003. Late Quaternary Palaeoecological Event Stratigraphy of the Horton Plains, Central Sri Lanka: with Contribution to the Recent Pollen Flora. Department of Physical Geography and Quaternary, Quaternary Geology 2, Stockholm University, Sweden. (Published Doctoral Thesis).
- Premathilake, R., Risberg, J., 2003. Late Quaternary history of the Horton Plains, central Sri Lanka. *Quaternary Science Review* 22, 1525-1541.
- Premathilake, R., 2006. Relationship of environmental changes in central Sri Lanka to possible prehistoric land-use and climate changes. *Palaeogeography, Palaeoclimatology, Palaeoecology* 240 (3/4), 468-496.
- Premathilake, R., 2012a. Late Pleistocene and early Holocene climate and environmental changes in the Horton Plains, central Sri Lanka. In: PAGES Focus 4 Biodiversity Theme Workshop: landscape planning for the future: using fossil records to map potential threats, opportunities and likely future developments for biodiversity and ecosystem services, University of Oxford, UK.
- Premathilake, R. 2012b. Human used montane ecosystem in the Horton Plains, central Sri Lanka - a link to glacial and early Holocene climate and environmental changes. *Quaternary Science Reviews* 50, 23-42.
- Premathilake, R., Gunathilake, A., 2013. Chronological framework of Asian southwest monsoon events and variations over the past 24,000 years in Sri Lanka and regional correlations. *Journal of National Science Foundation of Sri Lanka* 41, 3, 219-228.

939 Premathilake, R., 2015a. Late Pleistocene-Early Holocene Anthropogenic Environment from the  
 940 Horton Plains, Central Sri Lanka. In: Dissanayake, S., Rev. Chanaloka, P., Kodituwakku, N.,  
 941 (Eds.), Piyasatahan, Department of Archaeological Survey and Ministry of Cultural Affairs,  
 942 Colombo, pp. 118-156.

943 Premathilake, R., 2015b. Investigating the precursors and appearance of banana and rice  
 944 cultivation in Sri Lanka: with the background of long-term climate and environmental  
 945 changes. Association for Environmental Archaeology (AEA) Newsletter 128, 1-4.

946 Premathilake, R., 2016. Late Pleistocene human used *Musa* banana: phytoliths evidence from  
 947 Fahien rock shelter in Sri Lanka. Journal of Environmental Archaeology (AEA) Newsletter  
 948 133, 1-9.

949 Premathilake, R., Anupama, K., Rajan, K., Prasad, S., Orukaimani, G., Yathees Kumar, V.P.,  
 950 2017a. Implications of phytolith records from an Early Historic megalithic burial site at  
 951 Porunthal in Southern India. Journal of Archaeological Science: Reports 11, 491-506.

952 Premathilake, R., Kumara, A., Anupama, K., Pappu, S., Prasad, S., Gunnell, Y., Orukaimani, G.,  
 953 2017b. Phytoliths as indicators of Quaternary vegetation at the Palaeolithic sites of  
 954 Attirampakkam, India. Journal of Archaeological Science: Report.

955 Premathilake, R. and Hunt, C., 2018. Earliest *Musa* banana from the late Quaternary sequence at  
 956 Fahien Rock Shelter in Sri Lanka. Journal of Quaternary Science. In Press.

957 Rajagopalan, G., Sukumar, R., Ramesh, R., Pant, R.K., Rajgopalan, G., 1997. Late Quaternary  
 958 vegetal and climatic changes from tropical peats in southern India, an extended record up to  
 959 40,000 yr B.P. Current Science 73, 60-63.

960 Rajapaksha, U., 1998. Traditional food plants in Sri Lanka. Hector Kobbekaduwa Agrarian  
 961 Research and Training Institute, Colombo.

962 Reimer, P. J., Baillie, M. G. L., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Ramsey, C.  
 963 B., Buck, C. E., Burr, G. S., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P.,  
 964 Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kaiser, K. F., Kromer, B., McCormac, F.  
 965 G., Manning, S. W., Reimer, R. W., Richards, D. A., Southon, J. R., Talamo, S., Turney, C. S.  
 966 M., van der Plicht, J., Weyhenmeyer, C. E., 2009. INTCAL09 and MARINE09 radiocarbon  
 967 age calibration curves, 0-50,000 years cal. BP. Radiocarbon 51, 4, 1111-1150.

968 Ruddle K, Johnson D, Townsend P, Rees, J. 1978. Palm sago: A tropical Starch from Marginal  
 969 Lands. Melbourne: Australian National University Press.

970 Roach, M., 2013. The Bamboo Solution: Tough as steel, sturdier than concrete, full-size in a  
 971 year. Discover Magazine.

972 Roberts, P., Petraglia, M., 2015. Pleistocene rainforests: barriers or attractive environments for  
 973 early human foragers?, World Archaeology 47, 718-739.  
 974 <http://dx.doi.org/10.1080/00438243.2015.1073119>.

975 Roberts, P., Perera, N., Wedage, O., Deraniyagala, S., Perera, J., Eregama, S., Gledhill, A.,  
 976 Petraglia, M. D., Lee-Thorp, J. A., 2015. Direct evidence for human reliance on rainforest  
 977 resources in late Pleistocene Sri Lanka. Science 347, 1246-1249.

978 Roberts, P., Perera, N., Wedage, O., Deraniyagala, S., Perera, J., Eregama, S., Petraglia, M. D.,  
 979 Lee-Thorp, J.A., 2017. Fruits of the forest: Human stable isotope ecology and rainforest  
 980 adaptations in Late Pleistocene and Holocene (~36 to 3 ka) Sri Lanka. Journal of Human  
 981 Evolution 106, 102-118.

982 Saxena, A., Prasad, V., Singh, I. B., Chauhan, M.S., Hassan, R. 2006. On the Holocene record of  
 983 phytoliths of wild and cultivated rice from Ganga Plain: evidence for rice-based agriculture.  
 984 Current Science, 90, 11, 1547-52.

985 Simmonds, N. W., 1956. Botanical results of the banana collecting expedition, 1954-5. Kew  
986 Bulletin 11 (3), 463-490.

987 Simmonds, N. W., 1962. The Evolution of the Bananas. Tropical Science Series. Longmans,  
988 London.

989 Singh, I. B., 2005. Quaternary palaeoenvironments of the Ganga Plain and anthropogenic  
990 activity. Man and Environment 30, 1, 1-35.

991 Summerhayes, G. R., Leavesley, M., Fairbairn, A., Mandui, H., Field, J., Ford, A., Fullagar, R.,  
992 2010. Human Adaptation and plant use in Highland New Guinea 49,000 to 44,000 Years Ago.  
993 Science 330, 78-81.

994 Thompson, L.G., Yao, T., Mosley-Thompson, E., Davis, M.E., Henderson, K.A., Lin, P.- N.,  
995 2000. A high-resolution millennial record of the South Asian Monsoon from Himalayan ice  
996 cores. Science 289, 1916-1919.

997 Tsartsidou, G., Lev-Yadun, S., Albert, R.M., Miller-Rosen, A., Efstratiou, N., Weiner, S., 2007.  
998 The phytolith archaeological record: strengths and weaknesses evaluated based on a  
999 quantitative modern reference collection from Greece. Journal of Archaeological Science 34,  
1000 1262-1275.

1001 Vaughan, D. A. 1994. The Wild Relatives of Rice: A Genetic Resources Handbook. Manila:  
1002 International Rice Research Institute, Los Banos, Laguna (Philippines).

1003 Vaughan, D.A., Morishima, H., 2003. Biosystemetics of the Genus *Oryza*. In: Smith, C.W.,  
1004 Dilday, R. H., (Eds.), Rice origins, history, technology and production, John Wiley and Sons  
1005 Inc, New Jersey, pp.139-146.

1006 Vaughan, D.A., Lu, B.R., Tomooka, N. 2008. The evolving story of rice evolution. Plant  
1007 Science, 174, 394-408.

1008 Viet, N., 2001. Further studies of Hoabinhian in Vietnam, J. Southeast Asian Archaeology 21,  
1009 16-28.

1010 Vrydaghs, L., Ball, T., Volkaert, H., Houwe , I.V.D., Manwaring, J., De Langhe, E., 2009.  
1011 Differentiating the Volcaniform Phytoliths of Bananas: *Musa acuminata*. Ethnobotany  
1012 Research and Applications 7, 239-246.

1013 Weisskopf, A., Harvey, E., Kingwell-Banham, E., Kajale, M., Mohanty, R., Fuller, D. Q., 2014.  
1014 Archaeobotanical implications of phytolith assemblages from cultivated rice systems, wild  
1015 rice stands and macro-regional patterns. Journal of Archaeological Science 51,43-53.

1016 Wijeyapala, W. H., 1997. New Light on the Prehistory of Sri Lanka in the Context of Recent  
1017 Investigations of Cave Sites. Department of Archaeology, University of Peradeniya, Sri  
1018 Lanka. (Doctoral Thesis).

1019 Yang, Z., Barton, H. J., Wan, Z., Li, Q., Ma, Z., Li, M., Zhang, D., Wei, J., 2013. Sago-Type  
1020 Palms Were an Important Plant Food Prior to Rice in Southern Subtropical China. PLoS ONE  
1021 8(5), e63148. doi:10.1371/journal.pone.0063148.

1022 Zhijun, Z., 1998. The Middle Yangtze region in China is one place where rice was  
1023 domesticated: Dhvtolith evidence from the Diaotonghuan Cave, Northern Jiangxi. Antiquity  
1024 72, 885-97.

1025 Zong, Y., Chen, Z., Innes, J. B., Chen, C., Wang, Z., Wand, H., 2007. Fire and flood  
1026 management of coastal swamp enabled first rice paddy cultivation in east China. Nature, 449,  
1027 459-63.

1028 Zoysa Nde, Raheem, R., 1987. Sinharaja - a rainforest in Sri Lanka. March for Conservation,  
1029 Colombo.

1030 Zurro D, García-Granero J, lancelotti C, ET AL, 2016. Directions in current and future phytolith  
1031 research. Journal of Archaeological Science 68:112-117.  
1032

1033

1034     **Table 1**

1035

<b>Layers (Wijeyapala (1997))</b>	<b>Thick ness (m)</b>	<b>Contexts</b>	<b>Archaeological phases</b>	<b>Litho- stratigraphy</b>	<b>Colour</b>	<b>Biotic remains and archaeological finds</b>	<b>Cultural density</b>
Bedrock	NA	95	I	NA	NA	NA	NA
NA	0.40	92	II	Consolidated clast-rich loam	Yellowish brown	Ashy habitation deposits (hearths), human bone, wood charcoal, fragments of shells and fragments of microlithic flakes	Relatively low
L5	0.15	90, 91, 89	III	Moderately unconsolidated clast-rich loam	Pinkish grey to greyish brown	Ashy habitation deposits, charcoal, fragment of small mammals and human bones, burnt shells, fragments of <i>Canarium</i> nuts, hearths, microliths	Relatively high
L4	1.00	89, 88, 87	IV	Moderately unconsolidated clast-rich loam	Dark grey to brown	Ashy habitation deposits, charcoal, burnt shells, unburnt shell, human bones, <i>Canarium</i> nuts, microliths, red ochre, grindstones, postholes	High
NA		70	V				
L3	0.25	12/42/72	VI	Moderately unconsolidated loam	Brown	Charcoal rich habitation deposits, ashy, <i>Canarium</i> nuts, carnivore coprolites, bones, shells, unburnt shells, carnivores coprolites, wood, microliths. Fragmentary human skeleton found from the context 81, which has been directly dated to around 12,000 BP.	Relatively high
L3 NA		10, 11, 38, 48, 50, 62, 81	VI VI				

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045



1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062

**Table 2**

1063

Layers	Context (2010-2012 excavation)	Archaeological phase	Lab number	Lab age (BP)	$\delta^{13}\text{C}$	Calibrated dates BP (2 $\sigma$ )	Date range BC	Materials	Reference
Bedrock	95	I							Wijeyapala (1997)
	92	II	Beta-354918	43,030 $\pm$ 720	-27.0	44,952-47,854	45,905-43,003	Charred	Perera (2015)
L5	91	III	Beta-354920	34,610 $\pm$ 320	-25.7	38,477-39,852	37,903-36,528	Charred	Perera (2015)
	91	III	Beta-33294	33,070 $\pm$ 630	NA	35,768-38,716	36,767-33,819	Charcoal	Wijeyapala (1999)
	91	III	Beta-354914	32,890 $\pm$ 240	-26.9	36,263-37,844	35,895-34,314	Charred	Perera (2015)
L4	88	IV	Beta-33296	32,060 $\pm$ 630	NA	34,688-37,753	35,804-32,739	Charcoal	Wijeyapala (1997)
	87	IV	Beta-33299	30,060 $\pm$ 290	NA	33,674-34,644	32,695-31725	Charcoal	Wijeyapala (1997)
	87	IV	Beta-33295	24,470 $\pm$ 290	NA	27,860-29,089	27,140-25,911	Charcoal	Wijeyapala (1997)
NA	81	VI	UBA-19332	10,402 $\pm$ 45	-12.3	12,071-12,426	10,477-10,122	Freshwater shell	This paper
NA	81	VI	UBA-19331	10,344 $\pm$ 52	-29.7	11,991-12,402	10,453-10,042	Charred wood	This paper

1075

1076  
1077  
1078

1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087

## Figure and Table captions:

## Figure and Table captions for the main text:

### Figures

**Fig. 1. A:** Location of the Fahien rockshelter in Sri Lanka. Area marked by dark green color receives relatively high rainfall (>2,500 mm/yr) from the South West Monsoon while area marked by light green color receives relatively low (< 2,500 mm/yr) rainfall from the North East Monsoon. **B:** Contour map of the Rock Shelter.

**Fig. 2. X:** Plan of the rockshelter with excavated area. **Y:** Stratigraphy with the archaeological layers (L5-L3). Chronology of the Pleistocene sequence from the excavated area is shown. Contexts are marked by numbers. Locations of the monoliths and subsamples taken for phytolith analysis are indicated.

**Fig. 3.** Phytolith morphotypes from modern plants. Asteraceae (1-3): armed segmented hair cells with straight tips. Cyperaceae (4-10): 4-6: polyhedral conical projections, 7: polyhedral decorated cells, 8-9: three-four edge psilate-pitted (blocky), 10: epidermal cells. Panicoideae grass (11-15): 11: polylobate, 12-14: bilobate, 15: cross. Chloridoideae grass (16-19): saddle. Poideae grass (20-23): 20: elliptical psilate, 21-22: rectangle psilate, 23: round psilate. Bambusoideae (24-36): 24-31: rondel, 32-34: bulliform type B with angular-triangular flared keystone widely flared thin top symmetrical morphology. 35: elongate psilate. 36: large saddle. Poaceae (37-47): 37: elongate psilate, 38-39: elongate echinate, 40: elongate psilate, 41: bulliform type B with widely flared keystone thin top asymmetrical base morphology, 42-44: four edged smooth (blocky psilate), 45: double outline non-armed non-curved (trichome), 46-47: elongate pitted. Scale bar = 10  $\mu$ m.

**Fig. 4.** Phytolith morphotypes from modern plants. Moraceae/ *Artocarpus nobilis* (1-7), 1: double outline armed with acute tip attached to epidermal base, 2-4: double outline armed acute tip (trichome), 5-6: double outline armed acute tip attached to a round epidermal base with polyhedral cells, 7: epidermal polyhedral smooth cell. Bombacaceae/*Durio zibethius* (8-15): spheroidal-elliptical with nodular obtuse tip projections. Arecaceae/Palmae (15-21): 15: spheroidal-elliptical echinate from *Areca catchu*, 16-18: spheroidal echinate from *Borassus* sp., 19: spheroidal spinulose from *Caryota urans*, 20: spheroidal-elliptical spinulose from *Cocos nucifera*, 21: spheroidal echinate from *Phoenix zylanic*. 22: faceted from Moraceae/*Artocarpus nobilis*. 23-25: multi-faceted from Diterocarpaceae/*Shorea* sp. 26: Stomata from *Artocarpus nobilis*. Scale bar = 10  $\mu$ m.

**Fig. 5.** Phytolith morphotypes from modern plants. Seed phytoliths of *Musa accuminata* (1-9): 1-2: rectangular and square base with less protuberances, 3-6: irregular faceted and elongate with irregular short grooves, 7-9: faceted with nodule. Seed phytoliths of *Musa balbisiana* (10-17), 10-11: variable base with more protuberances. 12-15: irregular faceted with elongate well defined regular long grooves. 16-17: faceted with nodule. Volcaniform variants (V1-V8) from leaf of *Musa accuminata* (18) and *Musa balbisiana* (19). Crater, basal and protuberances of volcaniform are marked. Scale bar = 10  $\mu$ m. (cf. Ball et al., 2006).

**Fig. 6.** Phytolith morphotypes from modern rice plants. Bulliform type C phytoliths from domesticated rice leaves (1-7). 1: flared keystone moderately flared incurving sides, 2-4: flared keystone widely flared wide top asymmetrical base, 5: flared keystone moderately flared side not incurving, 6-7: keystone widely flared wide top with symmetrical base. 7: burnt bulliform. Diagnostic phytolith morphotypes of domesticated rice are marked with  $\psi$  symbol. 8: burnt scooped bilobate from ash sample (black color indicates burning), from traditional rice fields in Sri Lanka. 9-15: glume cell with various projections occurring on deeply serrated cells from traditional rice processing sites (e.g. named as *kamatha* in Sinhala language) in Sri Lanka. 16-17: glume cell with small projections occurring on deeply serrated cells from the archaeological context dated to 5/6<sup>th</sup> century BC and 12-13<sup>th</sup> century AD in southern India and Sri Lanka. Rice husks commonly produce these glume phytoliths. 18: large single-peaks, 19-20: double-peaks from domesticated rice husks. 21: double outlined non-armed with acute tip (trichome) from domesticated rice husks. 22: double outline non-armed curved small base with acute tips (trichome) from *Oryza* spp. 23: elongate cells with various projections (dendriforms) from *Oryza* spp. 24: epidermal non-curved striate phytoliths from domesticated rice husks. 25: large single and double peaks from *Oryza rufipogon* husks. 26: flared keystone moderately flared wide top symmetrical base from *Oryza rufipogon* leaves. 27-28: flared keystone moderately flared wide top with symmetrical base, and scooped bilobate and cross from *O. nivara* leaves. 29-30: large single peaks and 31: double peaks from *O. nivara* husks. Scale bar = 10  $\mu$ m.

**Fig. 7.** Phytolith percentages of the major economic plant taxa identified from the Rock Shelter samples analyzed.

**Fig. 8.** Wild banana phytoliths from Fahien Rock Shelter. 1-21: volcaniform variants (V1, V2, V3, V5 and V6) derived from wild banana leaves. 22-24: phytoliths from *Musa balbisiana* seeds. 25-26: phytoliths from *Musa accuminata* seeds; 27-28: small volcaniform phytoliths derived from wild banana seed. Scale bar = 10  $\mu$ m.

**Fig. 9.** Wild rice phytoliths from Fahien Rock Shelter. 1-5: flared keystone moderately flared wide top symmetrical base and flared keystone moderately flared incurving sides. Black color indicates burning. 6-14: glume cell with projections occurring on deeply serrated cells. 15-18: large single peaks. 19-22: large double peaks. Black color indicates burning. 23: glume cell with small projections. 24-25: double out lined non-armed curved small base with acute tips (trichome). 26: elongate echinate. 27-28: bilobate. Scale bar = 10  $\mu$ m.

**Fig. 10.** Phytoliths from Fahien Rock Shelter. Cyperaceae (1-6): 1-2: polyhedral base with conical projections (corn-shaped), 3-6: polyhedral decorated cells. Asteraceae (7): armed segmented hair cells with straight tips. Poaceae/Panicoideae (8): bilobate and polylobate. Poaceae/Chloridoideae (9-10): saddle. Bambusoideae (11-14): rondel. Arecaceae/Palmae (15-25): 15-18: spheroidal echinate, 19: spheroidal-elliptical spine, 20-21: spheroidal spinulose, 22-23: spheroidal-elliptical spinulose, 24-25: spheroidal spinulose. *Artocarpus* cf. *nobilis* (26-37): 26-32: double out lined armed with acute tip (trichome), 33-34: double outline armed acute tip attached round epidermal base, 35-37: round epidermal base comprised of polyhedral cells without arms. Bombacaceae/*Durio* cf. *zibethius* (38-40): spheroidal-elliptical nodular with obtuse

tip projections. Poaceae/grass (41-46): 41-43: bulliform types C, 44: elongate pitted, 45: elongate echinate, 46: double outlined non-armed with blunt tip. 47: large saddle from Bambusideae. 48: faceted. Burseraceae (49-50): faceted/scalloped elliptical (cf. Fig. 3 and 4). Scale bar = 10  $\mu$ m.

#### **Table captions:**

**Table. 1.** Summary of archaeology and stratigraphy described from the Fahien Rock Shelter (after Wijeyapala, 1997; excavation directed by Dr. Nimal Perera and Dr. Deraniyagala from the Department of Archaeology, Sri Lanka, 2010-2012). NA = Not available.

**Table. 2.** Chronology of the Fahien Rock Shelter (after Wijeyapala, 1997; Perera, 2015). NA = Not available.

#### **Supplementary Online Materials**

##### **Supplementary Tables:**

**Table S1.** Detailed stratigraphy of Fahien Rock Shelter

**Table S3:** Full phytolith raw data found from Fahien Rock Shelter

## **1. SOM 1**

### *1.1. Fossil phytolith extraction*

Sediment samples of 10-15 g were dried at 40° C and passed through 2 mm sieves. CaCO<sub>3</sub> was removed from the fraction passing the sieve using 10% HCl at 40° C. The material was centrifuged at 2000 rpm for 5 minutes. The supernatant was decanted and the remaining suspension checked with 1% AgNO<sub>3</sub> solution to ensure removal of CaCO<sub>3</sub>. The material was oxidized in 40 ml of 30% H<sub>2</sub>O<sub>2</sub> at 80-90° C in an oven for 2-3 hours. After cooling, a few drops of Ammonia solution were added to check for excess H<sub>2</sub>O<sub>2</sub>. The resulting suspension was passed through a 150  $\mu$ m sieve to remove coarse sand. The fraction less than 150  $\mu$ m was mixed with 20 ml 0.5% sodium hexametaphosphate. Clay particles were removed using density gradient techniques based on Stoke's Law. The silty fraction was removed and dried. 0.5 g of dry materials were mixed with 10 ml ZnBr<sub>2</sub> solution (density exactly adjusted to 2.35 gcm<sup>3</sup>) in a centrifuge tube. It was allowed to settle for 30 minutes and centrifuged at 2000 rpm for 30 minutes. The phytolith fraction was removed, mixed with 1N HCl and centrifuged at 2000 rpm for 5 minutes. The final phytolith fraction was mounted in Canada Balsam and observed at a magnification of x 400 under the Olympus BX51 microscope. Images critical to phytolith

identification were documented at a magnification of x 1000. Micrographs were taken using the F-View Soft Imaging System. Target counts were at least 250 (>250) and this was usually achieved. Counts of other siliceous microfossils (diatom) were also included. In order to understand the reliability of the archaeological phytolith assemblages in each sample and to trace vegetation signals, the relative abundance of each phytolith was calculated (Piperno, 1988). In this procedure, raw counts were transferred to percentage values (relative abundance) based on Tilia program. The basic sum included phytoliths derived from Poaceae. Other taxa were excluded from the sum. Phytolith diagrams were constructed using TILIA (Grimm, 1991-1993).

### *1.2. Modern phytoliths*

Modern samples (reference samples) from Sri Lanka and southern India were used. Taxonomic assignment of the reference samples were confirmed with the available Herbarium records at the Laboratory for Palaeoecology, Postgraduate Institute of Archaeology, French Institute of Pondicherry, India and Royal Botanic Garden, Kew (cf. Perera, 2017). Phytolith extraction from the samples was carried out both by ashing (Jenkins, 2009; Issaharou-Matchi et al., 2016) and chemical digenesis methods (Geiss, 1973 and Carter, 2007), because the results from those methodologies reflect the possible impact of such stresses on phytoliths at the field over a long period of time and during the laboratory processing as well. Plant samples of 25 g were cleaned using a bleaching agent (Alconox) and dried in a ceramic crucible at 50° C for five hours in order to remove adhered minerals. Subsamples of 1g of dry plant materials were taken for ashing and another subsample of 1 g was taken for chemical digenesis. In ashing method the materials were incinerated at 500° C for 5 hours in a ceramic crucible using a muffle furnace. Siliceous residues were further treated with 1N HCl, centrifuged at 2000 rpm for 5 minutes (centrifuging shall, hereafter, indicate this speed and time). The residual pellet was cleaned with distilled water and centrifuged. The pellet was treated with 5 ml of concentrated HNO<sub>3</sub> keeping in a water bath at 65° C for 2 hours and centrifuged. The pellet was cleaned with distilled water, centrifuged and then dried at 60° C. In the chemical digenesis method, the materials were treated with concentrated H<sub>2</sub>SO<sub>4</sub> (Sulphuric acid) at 80° C in a water bath for 4 hours until they dissolved. Then, the mixture was treated with 30% H<sub>2</sub>O<sub>2</sub> on a hot plate at 150° C for 2-3 hours till it turned colourless. The obtained residue was cleaned with distilled water and the pellet was air-dried. A

known weight of dry residue containing phytoliths obtained from both methods was separately mounted in both Canada balsam and Benzyl benzoate media.

## **2. SOM 2**

### *2.1. Phytolith taxonomy and modern plants*

Knowledge from the modern phytolith collections facilitated the fossil phytolith identification. In this collection, type slides were made from various plant parts (e.g., leaves, sheaf; seeds/husks, spikes, inflorescence/culm, roots and twigs) from a large number of traditional and modern cultivated and wild plants. This includes *Oryza nivara*, *O. rufipogon*, *Leersia* sp. *Oryza sativa*, *Panicum* sp., *Musa*-banana, *Artocarpus nobilis*, *Artocarpus* spp., *Durio zibethius*, bamboo, species from Asteraceae and Cyperaceae, Palmae, wild grasses, Dipterocarpaceae, Euphorbiaceae, Fabaceae, Verbenaceae and Burseraceae. The reference collection also contained phytolith type slides made from well preserved domesticated rice grains retrieved from archaeological contexts dating to the 6<sup>th</sup> and 3<sup>rd</sup> century BC and 12-13<sup>th</sup> century AD in southern India and Sri Lanka (Premathilake et al., 2017ab). To achieve additional knowledge on taxonomic identification, phytoliths were extracted from the samples obtained from several sites where traditional cereal processing activities (e.g. winnowing, de-husking and burning) was carried out in southern India and Sri Lanka. In our procedures, the most common morphological characteristics, general occurrence and appearance of key phytoliths, e.g. bulliforms, bilobates, saddles, rondel, elongate echinate, elongate sinuous, elongate projections, globular echinate, globular nodular spheroidal-elliptical nodular, double out line armed acute tips (trichome) double out line non-armed acute tip (trichome) and volcaniforms were comparatively studied (**Table S1**). The terminology was according to Madella et al. (2005), Pearsall and Dinan, 1992-2012 and Pearsall, 2015. Additional data from published works were used to differentiate taxa (e.g., Twiss et al., 1969; Bozarth et al. 1992; Pearsall and Dinan, 1992-2012; Fujiwara, 1993; Rapp and Mullholland, 1992; Mulholland, 1993; Pearsall et al., 1995; Houyuan et al., 1997; Zhijun, 1998; Zhao et al., 1998, Kealhofer and Penny, 1998; Runge, 1999; Zhao and Pearsall, 2000; Lu and Liu, 2000; Zheng et al., 2003; Thorn, 2004; Harvey and Fuller, 2005; Madella et al., 2005; Piperno, 2006; Saxena et al., 2006; Lejju et al., 2006; Ball et al., 2006, 2015; Bamford et al., 2006; Davey et al. 2007; Albert et al., 2009; Lu et al., 2009; Vrydaghs et al., 2003, 2009; Lentfer, 2009; Huang and Zhang, 2009; Mercader et al., 2010; Lippi et al., 2011; Weisskopf and Lee,

1277 2013; Weisskopf, 2014; Weisskopf et al., 2014, 2015ab; Yang et al., 2013; Lisztes-Szabó et al.,  
1278 2014; Zurro et al., 2016; Bates et al., 2017).

1279 **Table S1:** Phytoliths from modern taxa

<b>Taxa</b>	<b>Phytolith morphotypes</b>	<b>Botanical parts</b>
Banana (domesticated/wild)	Volcaniform (described in Table S4)	Leaf
Banana (wild)	Irregular faceted with elongated grooves	Seeds
<i>Artocarpus nobilis</i>	Double out line armed with acute tip (trichome)	Leaf
<i>Artocarpus nobilis</i>	Trichome base with polyhedral cells	Leaf
<i>Durio zibethius</i>	Globular-elliptical nodular obtuse tip projections (> 20µm)	Fruits and seeds
<i>Canarium zeylanicum</i> (Burseraceae)	Facetate/scalloped elliptical	Leaf
Burseraceae	Epidermal tissue polyhedral cells	Leaf
Wild rice ( <i>O. nivara/rufipogon</i> )	Small double-peak glume cell (25-30µm)	Husk
Wild rice ( <i>O. nivara/rufipogon</i> )	Large single-peak glume cell (20-30µm)	Husk
Wild rice ( <i>O. nivara/rufipogon</i> )	Small bumpy projections occurring on deeply serrated cells	Husk
Wild rice ( <i>O. nivara/rufipogon</i> )	Double out line non-armed with prominent base acute tip (trichome).	Husk and Inflorescence
Wild rice ( <i>O. nivara/rufipogon</i> )	Bulliform types-A. Flared keystone moderately flared deeply incurving side (1) and flared keystone widely flared wide top symmetrical base (2)	Leaf
Panicoid grass	Bilobate (rounded to square lobes)	Leaf and Inflorescence



Wild rice ( <i>O. nivara/rufipogon</i> )	Plate like epidermal (non-curvilinear with non-striate surface)	Husk
Pooid grass	Round to oblong	Leaf and Inflorescence
Pooid grass	Rectangle psilate	Leaf and Inflorescence
Pooid grass	Elliptic psilate	Leaf and Inflorescence
Pooid grass	Elongate sinuous	Leaf and Inflorescence
Chloridoid grass	Saddle	Leaf and Inflorescence
Poaceae (grass)	Plate like epidermal (non-curvilinear edge with smooth-pitted)	Husk
Poaceae (grass)	Elongate psilate and elongate potted	Leaf
Poaceae (grass)	Elongate echinate	Leaf
Poaceae (grass)	Elongate granulate	Leaf
Poaceae (willd grass)	Bulliform types-B: flared keystone slightly flared angular (1); flared keystone slightly flared rounded (2); flared keystone widely flared thin top symmetrical-asymmetrical base (3). Scale-like decorations < 9	Leaf
Pooid and Bambusoideae	Rondel	Leaf and trunk
Cyperaceae	Polyhedral decorated cell (achene)	Leaf
Cyperaceae	Polyhedral conical projection (hat shaped phytoliths)	Leaf
Cyperaceae	Blocky psilate-pitted	Leaf
Domesticated rice ( <i>O.</i>	Bulliform types-C: flared keystone	Leaf

<i>sativa</i> )	moderately flared weakly incurving side (1); flared keystone widely flared wide top asymmetrical base (2). Scale-like decorations > 9	
Domesticated rice ( <i>O. sativa</i> )	Scooped bilobate	Leaf and Inflorescence
Domesticated rice ( <i>O. sativa</i> )	Small double-peak glume cell (45-55 $\mu\text{m}$ )	Husk
Domesticated rice ( <i>O. sativa</i> )	Large single-peak glume cell (30-45 $\mu\text{m}$ )	Husk
Domesticated rice ( <i>O. sativa</i> )	Small bumpy projections occurring on deeply serrated cells	Husk
Domesticated rice ( <i>O. sativa</i> )	Double out line non-armed with acute tip attached to a weakly developed base (trichome).	Husk and Inflorescence
Domesticated rice ( <i>O. sativa</i> )	Bottle-like psilate and glob with short neck psilate	Husk
Domesticated rice ( <i>O. sativa</i> )	Plate like epidermal (non-curvilinear with striate surface)	Husk and Inflorescence
Asteraceae	Multicellular segmented hair cells	Leaf
Poaceae (wild grass)	Plate-like epidermal (non-curvilinear four edges with smooth-pitted) and blocky psilate bodies, double outline non-armed non-curved (trichome)	Leaf
Palmae	Globular echinate	Leaf and Inflorescence
Euphorbiaceae, Anacardiaceae and Annonaceae	Facetate	Leaf
Santalaceae and Zingiberaceae	Relatively large globular with projections	Leaf and Inflorescence

Euphorbiaceae, Anacardiaceae and Annonaceae	Sclereids	Leaf
Commelinaceae, Combretaceae, Cyperaceae, Poaceae and Fabaceae	Irregular shape blocky bodies with various projections	Leaf

### 2.1.2. Rice taxa identification

Comparative studies indicate that glume phytoliths i.e. double peak, single peak, elongate small round bumpy projection (glume cells), bumpy projections arising deeply serrated glume cells and elongate very deeply serrated sinuous edges glume cells classified into rice throughout the archaeological samples are broadly similar in to those from lemma, palea and rachilla i.e. seeds/husks samples from modern *Oryza nivara* and *O. rufipogon*. Bulliform phytoliths described as flared keystone moderately flared deeply incurving side (1) and flared keystone widely flared wide top symmetrical base (2) from modern *Oryza nivara* and *O. rufipogon* are similar to bulliform type A. The comparative studies indicate that on the lateral side of the bulliform type A from these wild rice species have less than 9 characteristic scale-like decorations (Table S1). These decorations are irregular and highly variable (cf. Wu et al., 2002). These species also produce regular short cell bilobate, epidermal non-curvilinear with non-striate surface and double out line non-armed with acute tip attached to a prominent base (trichome-1) phytoliths. The phytolith morphologies are consistent with the archaeobotanical evidence of rice. These key phytoliths are morphologically distinct from other wild rice i.e. *Leersia* sp. and wild grass populations studied. Many published works supported identification of rice (cf. Fujiwara, 1993; Harvey and Fuller, 2005; Huang and Zhang, 2009; Lippi et al., 2011; Tomber et al., 2011; Wang and Lu, 2012; Gu et al., 2013; Hu et al., 2013; Weisskopf, et al., 2014; Weisskopf et. al., 2015ab; Ball et al., 2016; Bates et al., 2017).

### 2.1.3. Wild *Musa banana* identification

Two groups of wild banana were recognized based on the morphology of seed phytoliths compared to the modern wild banana records (**Table S2**). In these groups, *Musa*-type 1 (crater width = 1-2.5µm; basal length = 8-10 µm; shape = rectangle and square base with less protuberances, and irregular faceted with elongate very irregular short grooves) and *Musa*-type 2 (crater width = 1-2.5µm; basal length = 3-7 µm; shape = variable base with more protuberances, and irregular faceted with elongate well defined regular long grooves) found in the late Pleistocene samples. The morphology of *Musa*-type 1 and 2 is closely similar to the morphology found from the modern seeds and leaves of *M. accuminata* and *M. balbisiana* respectively (**Table S2** and cf. Perera, 2017). However, based on morphometric evidence, leaf phytolith (volcaniform variants V1-V8) morphology of those two wild banana groups do not support the differentiation between *Musa*-type 1 and 2 (**Table S2**).

**Table S2:** Morphology of wild banana seed and leaf phytoliths. Total individual counts (N) per sample = 100

Modern wild banana seeds				
<i>M. accuminata</i>	1.97±0.30 range: 1-3.2	10.75±2.18 range: 8-18	Irregular faceted with elongate very irregular short grooves. Rectangle and square base with few verrucae.	
<i>M. balbisiana</i>	1.8±0.35 range: 1-3.2	8.62±2.25 range: 6-12	Irregular faceted with elongate well defined regular long grooves. Variable base with more verrucae.	
Modern wild banana leaf				
	Crater (µm)	Basal (µm)		
<i>M. accuminata</i>	7.07±0.89 range: 6.18-7.96	15.31±3.76 range: 11.55-9.07	Cone is visible with variants. V1 (84-85%), V3 (11-16%), V6 (1%), V5 (3-4%) are reported. V2, V4, V7, V8 are absent. Rectangle and square base with more protuberances dominate at the base of V6. Psilate-granulate with very few verrucate found.	Not available
<i>M. balbisiana</i>	7.62±1.06 range: 5.96-8.22	16.20±4.73 range: 11.46-21.24	Cone is visible with variants. V1 (67-85%), V3 (12-23%), V5 (1-12%), V6 (1-5), V8 (1-3), V4 (1) are reported. V2 and V7 are absent. Rectangle and square base with more protuberances dominate at the base of V6. Psilate-granulate with very few verrucate found.	Not available

1323

1324

1325

1326

1327

1328

1329

1330

1331

1332

1333

1334

Phytolith morphotypes and their affinities to botanical parts and plant taxa (cf. Table S1, S2 )			Analysed Sample					
Morphotype	Botanical parts	Taxa	C-81	C-87	C-88	C-89	C-91	C-92
Volcaniform	Banana leaves	Mucaceae/ <i>Musa</i> spp	18	31	130	31	60	65
Faceted	Banana seeds	Mucaceae/ <i>Musa</i> spp	1000	310	1173	1100	275	2620
Bulliform type-A	Rice leaves	<i>Oryza</i> sp.	6	1	15	2	5	18
Glume cell (single peak )	Rice husk	<i>Oryza</i> sp.		1	12	3	20	15
Glume cell (double peak)	Rice husk	<i>Oryza</i> sp.			15	3	7	25
Glume cell (deeply serrated sinuous edges)	Rice husk	<i>Oryza</i> sp.		2	25	3	29	42
Glume cell (deeply serrated sinuous/striate edges: plate-like)	Rice husk	<i>Oryza</i> sp.			12	1	2	2
Elongate dendritic	Rice husk/Inflorescence	<i>Oryza</i> sp.		1	6		7	7
Trichome (non-armed relatively round base acute tip)	Rice husk/Inflorescence	<i>Oryza</i> sp.			3		13	16
Glob with short neck psilate-granulate	Rice husk/Inflorescence	<i>Oryza</i> sp.					1	
Bottel-like psilate	Rice husk/Inflorescence	<i>Oryza</i> sp.						1
Bilobate convex ends	Leaves/Inflorescence	Panicoid		3				
Bilobate concave ends	Leaves/Inflorescence	Panicoid			5		2	5
Bilobate straight ends	Leaves/Inflorescence	Panicoid			2		2	3
Polylobate	Leaves/Inflorescence	Panicoid						2
Rounded psilate	Leaves/Inflorescence	Pooid	4	10	9	1	1	13
Elliptical psilate	Leaves	Pooid	3		1			3
Elongate sinuous	Leaves/Inflorescence	Pooid	5	15	30	2		2
Rectangle psilate	Leaves/Inflorescence	Pooid	150	25	300	100	100	209
Small saddle	Leaves/Inflorescence	Chloridoid			10		1	
Trichome (armed base not apparent straight edges acute tip)	Grass husk/Inflorescence	Poaceae/ (rice?)					5	1
Bulliform type-B	Grass leaves	Poaceae			3	3		5
Elongate deeply sinuous edges rounded bump	Grass/leaves Inflorescence	Poaceae			5	2		7
Elongate psilate	Grass leaves	Poaceae	25	10	45	2	10	45

Elongate wavy (echinate) symmetry	Grass leaves	Poaceae	1		5		20	13
Elongate wavy (echinate) asymmetry	Grass leaves	Poaceae			3			
Elongate echinate-baculate asymmetry	Grass leaves	Poaceae					5	2
Elongate granulate	Grass leaves	Poaceae	18		50		3	6
Rectangle-type granulate	Grass leaves	Poaceae	3	1	1	1	2	1
Elongate striate	Grass leaves	Poaceae			8	1		
Tabular trapizoid spine 60-70 µm	Grass leaves	Poaceae						3
Tabular trapizoid scrobiculate 60-70 µm	Grass leaves	Poaceae					1	
Undif.	Undif.	Poaceae	11	7	5	8	2	10
Large saddle	Leaves/steam	Bambusoideae	3		2			
Cone-like	Leaves/steam	Bambusoidea	36	12	20	15	12	17
Tabular trapizoid spine-scrobiculate 100-150 µm	Leaves	Dicot					1	
Globular-elliptical nodular obtuse tip projections	Seeds	<i>Durio</i> sp.	50	13	59	60		
Trichome (armed acute tip)	Leaves	<i>Artocarpus nobilis</i>	1	25	11	5	15	20
Trichome-base with polyhedral cell	Leaves	<i>Artocarpus nobilis</i>	1	4			3	10
Trichome (non-armed cone-like acute tip)	Leaves	Moraceae			3		15	23
Epidermal-type1: rectangle-square shaped/striate	Leaves	cf. <i>Artocarpus</i> sp.	3	3	10		1	25
Globular echinate >15 µm	Leaves	<i>Borass</i> spp	1	4	40	10		7
Globular echinate 7- 15 µm	Leaves/fruit	<i>Phonex</i> spp	100	90	230	300	31	160
Globular echinate < 7 µm	Leaves/steam	Palmae	16		11	50	10	50
Facetate (scalloped)	Undif.	Burseraceae	3					
Achene	Leaves	Cyperaceae			11		10	25
Globular psilate-verrucate	Undif.	Dicot		1	2		3	10
Globular irregular projection	Undif.	Dicot			6	1	10	20
Globular verrucate-baculate	Undif.	Dicot			1	1	1	2
Globular spine-baculate	Undif.	Woody dicot			4		1	5
Globular psilate 5 µm	Undif.	Dicot			1	1	2	2
Irregular facetate	Undif.	Woody dicot			4		9	12
Tabular facetate	Undif.	Dicot			2		10	5
Stomata	Undif.	Undif.					1	

Epidermal-type2 (hexagonal)	Undif.	Undif.			5		1	
Epidermal-type3 (honey comb)	Undif.	Undif.						1
Epidermal-type4 (elongate)	Leaves	Poaceae			2		1	8
Epidermal-type5 (Jigsaw-puzzle)	Leaves	Dicot			1		3	5
Undif. epidermal-types	Undif.	Undif.	1	1			1	6
Sclereids (tabular-like)	Undif.	Woody dicot					15	20
Blocky body perforate	Undif.	Dicot/monocot					2	3
Blocky body spine projection	Undif.	Dicot/monocot					5	2
Blocky body (irregular shaped) spinulose	Undif.	Dicot/monocot					3	2
Undif. (Dicot)	Undif.	Dicot	12		4		12	24
Tracheid	Undif.	Dicot/monocot			15	5		10
Burnt phytolith (bilobate/cross/bulliform/trichome/spherical echinate)	Leave/seed/husk	<i>Artocarpus/Musa/Oryza</i> spp/ <i>Palmae</i>	80	15	130	5	50	200
<i>Fallacia</i> sp	Diatom	Freshwater	2		2			
<i>Nitzschia</i> cf. <i>taylorii</i>	Diatom	Freshwater		1				3
<i>Surirella</i> sp.	Diatom	Freshwater						3
<i>Aulacoseria</i> cf. <i>granulata</i>	Diatom	Freshwater			7		2	2
<i>Melosira</i> cf. <i>lineata</i>	Diatom	Freshwater	2	4		7		2
<i>Navicula</i> cf. <i>leistikowii</i>	Diatom	Freshwater	200	10	5			
<i>Navicula</i> cf. <i>cryptotenella</i>	Diatom	Freshwater	20	1	9	2		3
<i>Navicula</i> sp.	Diatom	Freshwater	22		115	22	50	105
<i>Pinularia</i> spp	Diatom	Freshwater		1	7			4
<i>Eunotia</i> sp.	Diatom	Freshwater	1	1	4			
<i>Gomphonema</i> sp.	Diatom	Freshwater		1		1		
<i>Navicula</i> cf. <i>gregaria</i>	Diatom	Brackish/marine				1	1	1
<i>Achnanthes</i> cf. <i>hauckiana</i>	Diatom	Marine			30			10
<i>Amphora</i> cf. <i>coffeaeformis</i>	Diatom	Marine			1			

1335

1336 **Table. S3.** Detailed records of phytolith raw data found from the Fahien Rock Shelter.



1337

1338

## References

- Albert, R.M., Bamford, M.K., Cabanes, D., 2009. Palaeoecological significance of palms at Olduvai Gorge, Tanzania, based on phytolith remains. *Quaternary International* 193, 41-48.
- Ball, T., Vrydaghs, L., Houwe, I.V.D., Manwaring, J., De Langhe, E., 2006. Differentiating banana phytoliths, wild and edible: *Musa acuminata* and *Musa balbisiana*. *Journal of Archaeological Science* 33, 1228-1236.
- Ball, T., Chandler-Ezell K., Dickau, R., Duncan, N., Hart, T. C., Iriarte, J., Lentfer, C., Logan, A., Lu, H., Madella, M., Pearsall, D. M., Piperno, D. R., Rosen, A. M., Vrydaghs, L., Weisskopf A., Zhang, J., 2016. Phytoliths as a tool for investigations of agricultural origins and dispersals around the world. *Journal of Archaeological Science* 68, 32-45.  
<http://dx.doi.org/10.1016/j.jas.2015.08.010>.
- Bamford, M. K., Albert, R. M., Cabanes, D., 2006. Plio-Pleistocene macroplant fossil remains and phytoliths from Lowermost Bed II in the eastern palaeolake margin of Olduvai Gorge, Tanzania. *Quaternary International* 148, 95-112.
- Bates J, Singh, R. N., Petrie, C. A., 2017. Exploring Indus crop processing: combining phytolith and macrobotanical analyses to consider the organization of agriculture in northwest India c. 3200-1500 BC. *Vegetation History Archaeobotany* 26, 25-41. DOI 10.1007/s00334-016-0576-9.
- Bozarth, S. R., Rapp, G. R., Mulholland, S. C., 1992. Classification of opal phytoliths formed in selected dicotyledons native to the Great Plains. *Phytolith Systematics*. Plenum Press, New York.
- Carter, J. A., 2007. Ancient Climate and Environmental History from Phytoliths-occluded Carbon. Doctoral Thesis, Department of Geology, Victoria University of Wellington.
- Davey, F.I., Taylor, D., Dodson, J., Atahan, P., Zheng, H., 2007. Wild and domesticated form of rice (*Oryza* sp.) in early agriculture at Qingpu, lower Yangtze, China: evidence from phytolith. *Journal of Archaeological Science* 34, 2101-2108.
- Fujiwara, H., 1993. Research into the history of rice cultivation using plant opal analysis. In: Pearsall, D.M., Piperno, D.R. (Eds.), *Current Research in Phytolith Analysis: Applications in Archaeology and Palaeoecology*, MASCA Research Papers in Science and Archaeology. Vol. 10. University of Pennsylvania, Philadelphia, pp. 147-158.
- Geiss, J. W., 1973. Biogenic silica in selected species of deciduous angiosperms. *Soil Science* 116, 113-119.
- Grimm, E. C., 1991-1993. TILIA 2.00. Illinois State Museum Research & Collections Centre, 1920 South 101 2 St., Springfield, IL 62703, USA.
- Harvey, E. L., Fuller, D. Q., 2005. Investigating crop processing using phytolith analysis: the example of rice and millets. *Journal of Archaeological Science* 32, 739-752.
- Houyuan, L. Naiqin, W., Baozhu, L., 1997. Recognition of rice phytoliths. In: Pinilla, A., Juan-Tresserras, J., Machado, M. J., (Eds.), *The state-of-the art of phytoliths in soil and plant*, Monografias del Centro de Ciencias Medioambientales, Consejo superior de Investigaciones científicas, Madrid, pp. 159-174.
- Huang, F., Zhang, M., 2009. Pollen and phytolith evidence for rice cultivation during the Neolithic at Longqiuzhuang, eastern Jianghuai, China. *Vegetation History Archaeobotany* 18, 161-168.
- Hu, L., Chao, Z., Gu, M., Li, F., Chen, L., Liu, B., Li, X., Huang, Z., Li, Y., Xing, B., Dai, J., 2013. Evidence for a Neolithic Age fire-irrigation paddy cultivation system in the lower Yangtze River Delta, China. *Journal of Archaeological Science* 40, 72-78.

1385 Issaharou-Matchi, I., Barboni, D., Meunier, J. D., Saadou, M., Dussouillez, P., Contoux, C.,  
1386 Zirihi-Guede, N., 2016. Intraspecific biogenic silica variations in the grass species *Pennisetum*  
1387 *pedicellatum* along an evapotranspiration gradient in South Niger. *Flora - Morphology,*  
1388 *distribution, Functional Ecology of Plants*, 220, 84-93.

1389 Jenkins, E., 2009. Phytolith taphonomy: a comparison of dry ashing and acid extraction on the  
1390 breakdown of conjoined phytoliths formed in *Triticum durum*. *Journal of Archaeological*  
1391 *Science* 36, 2402–2407.

1392 Kealhofer, L., Penny, D., 1998. A combine phytolith and pollen records for fourteen thousand  
1393 years of vegetation change in Northeastern Thailand. *Review of Palaeobotany and Palynology*  
1394 103, 83-93.

1395 Lejju, J. B., Robertshaw, P., Taylor, D., 2006. Africa's earliest bananas? *Journal of*  
1396 *Archaeological Science* 33, 102-113.

1397 Lentfer, C.J., 2009. Tracing Domestication and Cultivation of Bananas from Phytoliths: An  
1398 update from Papua New Guinea. *Ethnobotany Research and Applications* 7, 247-270.

1399 Lippi, M. M., Gonnelli, T., Pallecchi, P., 2011. Rice chaff in ceramics from the archaeological  
1400 site of Sumhuram (Dhofar, Southern Oman). *Journal of Archaeological Science* 38, 1173-  
1401 1179.

1402 Lisztes-Szabó, Z., Kovacs, S., Peto, A., 2014. Phytolith analysis of *Poa pratensis* (Poaceae)  
1403 leaves. *Turkish Journal of Botany*, 38, 851-863.

1404 Lu, H., Liu, K., 2000. Illustrations of phytoliths from grasses in China and the USA. Institute of  
1405 Geology and Geophysics. Chinese Academy of Sciences, Beijing.

1406 Lu, H., Zhang, J., Liu, K., Wu, N., Li, Y., Zhou, K., Ye, M., Zhang, T., Zhang, H., Yang, X., Shen,  
1407 L., Xu, D., Li, Q., 2009. Earliest domestication of common millet (*Panicum miliaceum*) in  
1408 East Asia extended to 10,000 years ago. *PNAS*, 106, 7367-7372.

1409 Madella M, Alexandre A, Ball T (2005). International Code for phytolith nomenclature 1.0  
1410 *Annals of Botany* 96, 253-260.

1411 Mercader, J., Astudillo, F., Barkworth, M., Bennett, T., Esselmont, C., Kinyanjui, R., Grossman,  
1412 D. L., Simpson, S., Walde, D., 2010. Poaceae phytoliths from the Niassa Rift, Mozambique  
1413 *Journal of Archaeological Science* 37, 1953-1967.

1414 Mulholland, S. C., 1993. A test of phytolith analysis at Big Hidatsa, North Dakota. In: Pearsall,  
1415 D. M. and Piperno, D. R. (Eds.), *Current research in phytolith analysis: application in*  
1416 *archaeology and palaeoecology. MASCA research papers in science and archaeology* 10, The  
1417 University of Pennsylvania, Philadelphia, pp. 131-145.

1418 Pearsall, D. M., Dinan, E., 1992-2012. University of Missouri phytolith classification system. In:  
1419 *Studies in Phytolith Systematics*, edited by G. Rapp, Jr. and S. Mulholland, Plenum Press,  
1420 New York, pp. 37-64.

1421 Pearsall, D. M., Piperno, D. R., Dinan, E. H., Umlauf, Z., Benfer Jr., R. A., 1995.  
1422 Distinguishing rice (*Oryza sativa* Poaceae) from wild *Oryza* species through phytolith  
1423 analysis. Results of preliminary research. *Economic Botany* 49, 183-186.

1424 Pearsall, D. M., 2015. Phytolith Classification System: Notes and Guidelines for Use.  
1425 Department of Anthropology, University of Missouri, Columbia.

1426 Perera, P., 2017. A record of modern phytoliths morphology from *Musa* bananas: an approach to  
1427 identify banana phytoliths on archaeological context in Sri Lanka. Postgraduate Institute of  
1428 Archaeology, University of Kelaniya. (Master of Philosophy Thesis).

1429 Piperno, D. R., 1988. Phytolith analysis: An archaeological and geological perspective. San  
1430 Diego: Academic Press.

- Piperno, D. R., 2006. *Phytoliths: a comprehensive guide for archaeologist and paleoecologist*. Alta Mira press, Rowman and Littlefield Publishers, Inc. UK.
- Premathilake, R., Anupama, K., Rajan, K., Prasad, S., Orukaimani, G., Yathees Kumar, V. P., 2017a. Implications of phytolith records from an Early Historic megalithic burial site at Porunthal in Southern India. *Journal of Archaeological Science Reports* 11, 491-506.
- Premathilake, R., Akhilesh, K., Anupama, K., Pappu, S., Prasad, S., Gunnell, Y., Orukaimani, G., 2017b. Phytoliths as indicators of Quaternary vegetation at the Paleolithic site of Attirampakkam, India. *Journal of Archaeological Science Reports* 14, 470-490.
- Rapp Jr., G., Mulholland, S.C., (Eds.), 1992. *Phytolith Systematics*. Plenum Press, New York.
- Runge, F., 1999. The opael phytolith inventory of soil in central Africa-Quantities, shapes, classification and spectra. *Review of Palaeobotany and Palynology* 107, 23-53.
- Wu, N., Berne, S., Saito, Y., Liu, B., Wang, L., Lu, H., Liu, Z., 2002. Rice domestication and climatic change: phytolith evidence from East China. *Boreas* 31, 378-385.
- Thorn, V. C., 2004. Phytolith evidence for C<sub>4</sub>-dominated grassland since the early Holocene at Long Pocket, northeast Queensland, Australia. *Quaternary Research* 61, 168-180.
- Tomber, R., Cartwright, C., Gupta, S., 2011. Rice temper: technological solutions and source identification in the Indian Ocean. *Journal of Archaeological Science* 38, 360-366.
- Twiss, P.C., Suess, E., Smith, R.M., 1969. Morphological classification of grass phytoliths. *Soil Science Society of America Proceedings* 33, 1, 109-115.
- Saxena, A., Prasad, V., Singh, I. B., Chauhan, M.S., Hassan, R. 2006. On the Holocene record of phytoliths of wild and cultivated rice from Ganga Plain: evidence for rice-based agriculture. *Current Science*, 90, 11, 1547-52.
- Vrydaghs, L., Swennen, R.R., Mbida, C., Doutrelepon, H., De Langhe, E., De Maret, P., 2003. The banana phytolith as direct marker of early agriculture. In: Hart, D.M., Wallis, L.A.A. (Eds.), *A review of the evidence of phytolith and starch research in the Australian-Pacific-Asian regions: the state of the art*. Terra Australis 19. Pandanus Books, Research School of Pacific and Asian Studies, Australian National University, Canberra, pp. 17-185.
- Wang, C., Lu, H., 2012. Research progress of fan shaped phytoliths of rice and relevant issues. *Quaternary Science* 32, 269-281.
- Gu, Y., Zhao, Z., Pearsall, D. M., 2013. Phytolith morphology research on wild and domesticated rice species in East Asia. *Quaternary International* 287, 141-148.
- Weisskopf, A. R., Lee, G.A., 2013. Phytolith identification criteria for foxtail and broomcorn millets: a new approach to calculating crop ratios. *Archaeological Anthropological Science*. DOI 10.1007/s12520-014-0190-7.
- Weisskopf A., Harvey E., Kingwell-Banham E., Kajale M., Mohanty R., Fuller DQ (2014). Archaeobotanical implications of phytolith assemblages from cultivated rice systems, wild rice stands and macro-regional patterns. *Journal of Archaeological Science* 51, 43-53.
- Weisskopf AR (2014). Millets, rice and farmers: phytoliths as indicators of agricultural, social and ecological change in Neolithic and Bronze Age Central China. *Bar International Series* 2589. Archaeopress, Oxford.
- Weisskopf AR, Deng Z, Qin L, Fuller D (2015a). The interplay of millets and rice in Neolithic central China: Integrating phytoliths into the archaeobotany of Baligang. *Archaeological Research in Asia* 4, 36-45.
- Weisskopf, A. R., Qin, L., Ding, J., Ding, P., Sun, G., Fuller, D. Q., 2015b. Phytoliths and rice: from wet to dry and back again in the Neolithic Lower Yangtze. *Antiquity* 89, 347, 1051-1063.

- Yang, Z., Barton, H. J., Wan, Z., Li, Q., Ma, Z., Li, M., Zhang, D., Wei, J., 2013. Sago-Type Palms Were an Important Plant Food Prior to Rice in Southern Subtropical China. PLoS ONE 8(5), e63148. doi:10.1371/journal.pone.0063148.
- Zhao Z., 2001. New Archaeobotanic Data for the Study of the Origins of Agriculture in China. Current Anthropology 52 (S4), S295-S306.
- Zhao, Z., Pearsall, D. M., 2000. Late Pleistocene/Holocene environments in the middle Yangtze River Valley, China and rice (*Oryza sativa* L.) domestication: The phytolith evidence. Geoarchaeology 15, 203-222.
- Zheng, Y., Dong, Y., Matsui, A, Udatsu, T., Fujiwara, H., 2003. Molecular genetic basis of determining subspecies of ancient rice using the shape of phytoliths. Journal of Archaeological Science 30, 1215-1221.
- Zhijun, Z., 1998. The Middle Yangtze region in China is one place where rice was domesticated: Phytolith evidence from the Diaotonghuan Cave, Northern Jiangxi. Antiquity 72, 885-97.
- Zurro, D., García-Granero, J., lancelotti, C., Madella, M., 2016. Directions in current and future phytolith research. Journal of Archaeological Science 68, 112-117.





Fig. 1




**Fig. 2**






**Fig. 3.**






**Fig. 4.**






**Fig.5.**






**Fig.6**



**Fig. 7.**





**Fig. 8**






Fig. 9






Fig. 10